

1932, No. 4

DECEMBER 31

COPEIA

A JOURNAL OF COLD BLOODED
VERTEBRATES

Established in 1913

PUBLISHED BY
THE AMERICAN SOCIETY OF ICHTHYOLOGISTS
AND HERPETOLOGISTS

CONTENTS

BARTON WARREN EVERMANN, 1853-1932. <i>By G. D. Hanna</i>	161
REPORT ON TWO COLLECTIONS OF LOWER CALIFORNIA MARINE FISHES. <i>By Joseph H. Wales</i>	163
FEEDING HABITS AND MOLT OF <i>CROTALUS CONFLUENTUS OREGANUS</i> IN CAPTIVITY. <i>By Tracy I. Storer and Beryl M. Wilson</i>	169
BLANDING'S TURTLE, <i>EMYS BLANDINGII</i> (HOLBROOK) IN PENNSYLVANIA. <i>By M. Graham Netting</i>	173
HERPETOLOGICAL RECORDS AND NOTES FROM THE VICINITY OF TUCSON, ARIZONA, JULY AND AUGUST, 1930. <i>By F. Willis King</i> ..	175
HERPETOLOGICAL NOTES—The Status of Two Peruvian Lizards, <i>by</i> <i>H. W. Parker</i> : p. 178.—Comments on Polypedatidae, <i>by Benjamin</i> <i>B. Leavitt</i> : p. 179.—An Aberrant <i>Natrix clarkii</i> (Baird and Girard), <i>by Morrow J. Allen</i> : p. 179.	
ICHTHYOLOGICAL NOTES—Fish Notes for 1931 and 1932 from Sandy Hook Bay, <i>by C. M. Breder, Jr.</i> : p. 180.—The Status of <i>Osphronemus</i> <i>saigonensis</i> Borodin, <i>by Hugh M. Smith</i> : p. 180.—A New Record of <i>Ostracion diaphanum</i> Bloch and Schneider from Cali- fornia, <i>by Howard R. Hill</i> : p. 181.—A Record of the <i>Luvaru</i> (<i>Luvarus imperialis</i> Rafinesque) from Southern California, <i>by</i> <i>Howard R. Hill</i> : p. 181.—A Case of Arterial Constriction of the Spleen in the Dogfish, <i>by Charles E. Burt</i> : p. 181.—Blue Perch, <i>by J. R. Dymond</i> : p. 182.	
REVIEWS AND COMMENTS—Progress on the Bashford Dean Memorial Volume on Archaic Fishes, <i>by William K. Gregory</i> : p. 182.	
EDITORIAL NOTES AND NEWS—Piscium Catalogus: p. 184.—New York Zoological Society: p. 184.—Herpetological Items: p. 184.—1933 Meeting: p. 185.—National Museum: p. 185.—The National Aquarium Society: p. 185.—The Plural of Fish: p. 185.—Recent Death, Dr. William Patten: p. 185.	
INDEX FOR 1932.....	186

51

53

59

73

85

4

D

E

3

1

3

2

MI

1932, No. 4

DECEMBER 31

COPEIA

A JOURNAL OF COLD BLOODED
VERTEBRATES

Established in 1913

PUBLISHED BY
THE AMERICAN SOCIETY OF ICHTHYOLOGISTS
AND HERPETOLOGISTS

Barton Warren Evermann, 1853-1932

IN the death of Dr. Evermann biological science lost one of its most enthusiastic supporters, and American ichthyology was an especially heavy loser. Within a comparatively few years there have passed David Starr Jordan, Charles H. Gilbert, Carl H. Eigenmann and Barton Warren Evermann, four zoologists who dominated the study of fishes in this country for half a century and to such an extent that their combined efforts may come to be known as a "school" or a "regime." They approached many of their subjects in a similar manner, that is from the side of taxonomy. One of the characteristics of the "school" was the frequency with which their papers appeared under joint authorship.

It seemed to those of us who were closely associated with Dr. Evermann that he began to decline in 1923 when, on a trip to the lower Colorado River, he contracted a severe bronchial disturbance which never thereafter completely vanished. With each succeeding attack he was not able to completely regain the loss. Nevertheless, an indomitable will to keep doing things, and a physique which was normally very robust, kept him at active duty until June 21, 1932. He then went to Stanford Hospital for a brief rest. One complication followed another and he finally succumbed to pneumonia on September 27, 1932.

Dr. Evermann was born in Iowa on October 24, 1853, and was educated in Indiana, the state he always considered home. The friendship which began in his college days with Dr. Jordan, then president of Indiana University, was very intimate and lasting. It resulted in a long list of papers and books on fishes under joint authorship. This mutual willingness to help each other was not confined to joint works but extended to real assistance in the preparation of their own papers. They seemed to take it as a matter of course that they should read and correct each other's proof sheets and supply illustrations and references.

Dr. Evermann left a card index of titles to his publications, the last number being 387. His bibliography was published in 1905 (*Indiana University Bulletin*, vol. 2, no. 6, March 1905, pp. 84-90), and it is hoped that the remaining titles can be brought up to date. All of the latter ones (1920-1930) may be found in the various annual reports of the Director of the California Academy of Sciences. The last publication prior to his death was a review of W. S. Blatchley's book "My Nature Nook" (*Science*, vol. 76, no. 1959, July 15, 1932, pp. 57-58). He left finished and ready for the press a paper on the fishes of

the Revillagigedo Islands Expedition of 1925 (joint authorship with H. W. Clark) and his annual report as Director of the Academy for 1931. Plans had been drawn for several other publications but so far as known he had not begun the actual preparation of any of them. It is seldom that so little unfinished work is left behind.

The fame of the man rests largely on his studies of fishes but an analysis of his card index shows that he published widely on other branches. There are, as a very general classification, 196 papers on fishes; 59 on birds; 30 on mammals; and the remainder on various subjects. Many of his undertakings were of monographic nature, and involved an enormous amount of routine and detail. Most of his work was done at the same time the duties of a strenuous administrative position were being fulfilled. He attended to a voluminous correspondence, largely by rapid fire dictation, but his manuscripts were laboriously and studiously written in long hand.

Dr. Evermann taught in the public schools of Indiana and California for 10 years after leaving college and this experience left a lasting impression upon him; at heart he was a teacher. He held various positions with the U. S. Bureau of Fisheries from 1886 to 1914, his most prolific period as a writer. He went to the California Academy of Sciences in 1914 as Director, a position which he held until his death. During his regime many advances were made by the Academy, the most important perhaps being the establishment of the Steinhart Aquarium and the acquisition of the great collection of fishes formerly belonging to Indiana University. More than 20 volumes of scientific reports were published under his direction and expeditions were dispatched far and wide.

Many species of animals and plants and five genera of fishes have been named for him. The genera are: *Evermannia* Jordan, 1895; *Evermanella* Fowler, 1901; *Evermanella* Eigenmann, 1903 (changed to *Evermanolus* by Eigenmann in 1907); and *Evermannichthys* Metzelaar, 1919. The name Mt. Evermann for the highest peak of Socorro Island has been accepted by the Mexican Government and by the U. S. Geographic Board.

On October 24, 1875, Dr. Evermann married Meade Hawkins at Burlington, Indiana; she died at Berkeley, California, on February 9, 1929. A son and a daughter survive: Toxaway Bronte and Mrs. Edith Humphrey, both of Washington, D.C. In addition to these, a sister, Miss Dora Evermann was with him when he died. His body was cremated and the ashes were buried in Burlington Cemetery at Burlington, Indiana, on October 22, 1932.

G. D. HANNA

*California Academy of Sciences,
San Francisco, California.*

Report on Two Collections of Lower California Marine Fishes

By JOSEPH H. WALES

SIXTY of the sixty-nine species here reported were collected by the writer while he was a member of the California Division of Fish and Game Lower California Expedition of 1931. This boat trip was made to La Paz in the Gulf of California and back between March 31 and April 17.

The fish were collected by seining, by hook and line, by dip-netting under a drop-light at night, and in tide pools. All of the methods were fairly successful.

Ten species were collected by Ira L. Wiggins while botanizing in Lower California during April and May, 1931, and through his courtesy are included in this report.

The collections are in the Natural History Museum, Stanford University.

None of the species here included are new but the ranges of at least sixteen have been extended.

The writer wishes to thank the California Division of Fish and Game for permitting him to accompany the expedition. H. R. Hill of the Los Angeles Museum, kindly assisted in the identification of *Mycteroperca* and *Lutjanus*.



Map of Lower California, locating points mentioned in text.

Galeidae—Requiem Sharks

1. *Triakis semifasciata* Girard.—Ensenada Bay, April 1, one specimen three feet long, seined. Turtle Bay, April 4, many seined.

Rhinobatidae—Guitar Fishes

2. *Rhinobatos productus* Ayres.—Turtle Bay, April 4, many seined. Magdalena Bay, April 7, four seined.

Dasyatidae—Sting Rays

3. *Dasyatis dipterura* Jordan and Gilbert.—La Paz, April 10, one seined, 375 mm broad.

Myliobatidae—Eagle Rays

4. *Myliobatis californicus* Gill.—Turtle Bay, April 4, five seined.

Myridae—Worm Eels

5. *Myrophis vafer* Jordan and Gilbert.—Magdalena Bay, April 7, two specimens, 170 mm and 130 mm long. La Paz, April 9, three specimens, 260 mm, 280 mm and 300 mm. All five specimens were taken by drop light. All had insertion of dorsal fin slightly closer gill slit than anus. The Magdalena specimens seem to constitute a northern record for the west side of the peninsula.

Muraenidae—Morays

6. *Gymnothorax dovii* (Günther).—La Paz, April 10, one specimen 480 mm long, from tide pool. Head 2.5 in trunk.

Albulidae—Bone Fishes

7. *Albula vulpes* (Linnaeus).—Turtle Bay, April 4, fourteen specimens, 35-70 mm, seined.

Dussumieriidae—Round Herrings

8. *Etrumeus micropus* Temminck and Schlegel.—Turtle Bay, April 4, one seined.

Clupeidae—Herrings

9. *Sardinops caerulea* (Girard).—Turtle Bay, April 4, many seined. Magdalena Bay, April 7, many seined. La Paz, April 9, one specimen 42 mm long, by drop-light.

Engraulididae—Anchovies

10. *Cetengraulis mysticetus* (Günther).—Magdalena Bay, April 6, six specimens about 135 mm. long, seined by tuna boat the previous night in the bay. Apparently the first record for the west side of the peninsula.

11. *Engraulis mordax mordax* Girard.—Turtle Bay, April 4, one specimen 70 mm, seined. La Paz, April 9, one taken by drop-light. Apparently the first record for the Gulf.

Synodontidae—Lizard Fishes

12. *Synodus*, sp. Mainland opposite Cerralbo Island, April 11, one post-larval specimen 60 mm long, by drop-light. Transparent with nine patches of melanophores on dorsal surface from head to tail; ten round black spots situated internally along the median ventral axis of the body, the first two pairs separate, the rest connected; base of caudal black.

Belonidae—Needle Fishes

13. *Strongylura exilis* (Girard).—Turtle Bay, April 4, five specimens, 500-650 mm, seined.

Hemirhamphidae—Half Beaks

14. *Hyporhamphus roberti* (Cuvier and Valenciennes).—Turtle Bay, April 4, six specimens, 110-135 mm, seined. La Paz, April 10, three specimens about 270 mm, by seine and drop-light.

Exocoetidae—Flying Fishes

15. *Fodiator acutus* (Cuvier and Valenciennes).—La Paz, April 9, four specimens, 165-185 mm, by drop-light.

Atherinidae—Silversides

16. *Atherinopsis sonorae* Osburn and Nichols.—Turtle Bay, April 4, hook and line.

17. *Atherinops affinis magdalenae* Fowler.—Magdalena Bay, April 7, many seined.

Mugilidae—Mulletts

18. *Mugil curema* Cuvier and Valenciennes.—Turtle Bay, April 4, many seined. Magdalena Bay, April 7, many seined. La Paz, April 9, many seined.

19.—*Mugil cephalus* Linnaeus.—Valencia Beach, April 2, one specimen collected by Ira L. Wiggins. San Jose del Cabo, May 9 and May 14, five and six young specimens respectively, collected by Ira L. Wiggins.

20. *Chaenomugil proboscideus* (Günther).—Near east side of peninsula opposite Cerralbo I., April 11, one specimen by drop-light.

Sphyraenidae—Barracudas

21. *Sphyraena argentea* Girard.—Magdalena Bay, April 8, two specimens about 370 mm, hook and line.

Scombridae—Mackerels

22. *Scomberomorus sierra* Jordan and Starks.—La Paz, mouth of south channel, April 9, one 750 mm specimen, hook and line.

23. *Pneumatophorus diego* (Ayres).—Magdalena Bay, April 7, many seined. Apparently the first Lower California record.

24. *Sarda chilensis* Cuvier and Valenciennes.—Turtle Bay, April 4, many seined.

25. *Katsuwonus pelamis* (Linnaeus).—Fifty miles south of Santa Margarita Island, April 8, three specimens, hook and line.

26. *Katsuwonus lineatus* Kishinouye.—Off Santa Margarita Island, April 8, two specimens about 580 mm long. These two apparently constitute the third record of the species. The second record was by G. C. Thomas, Jr., and G. C. Thomas, III (Game Fish of the Pacific, Southern Californian and Mexican, 1930, pl. opposite p. 28).

27. *Neothunnus macropterus* (Schlegel).—La Paz, south channel, April 9, one specimen, 600 mm long. Head 3 in body, eye 5.5 in head, maxillary reaching pupil, pectoral 7/8 head, ventrals 2.8 head, anal lobe length of ventrals, lobe of soft dorsal slightly longer, longest dorsal spine 2.5 in head, caudal lobes 1.4 head. Dorsal XIII, 13, 9 finlets. A. 11, 8 finlets.

Carangidae—Pampanos

28. *Oligoplites mundus* Jordan and Starks.—La Paz, April 10, one specimen seined.

29. *Trachurops crumenophthalmus* (Bloch).—La Paz, April 10, one specimen seined.

Serranidae—Sea Basses

30. *Mycteroperca xenarcha* Jordan.—Between Seal Rock and Cerralbo Island, April 11, one specimen, 775 mm long. Anal III, 12; 4th anal ray 2 in head; 10th dorsal ray 3.67 in head; 3rd dorsal spine longest, 4 in head;

interorbital width 4.5 in head; 2 canines on each side of upper jaw; 1 on each side of lower jaw; maxillary 2.2 in head. Color dark brown, heavily speckled on head, body, and fins with darker spots smaller than pupil; spots showing up on lighter areas such as ventral part of body and opercles; fins all dark brown with spots. This specimen has been compared with one of *xenarcha* in the Stanford collection, and found to agree reasonably well except that the Stanford specimen does not show the black spots.

31. *Paralabrax nebulifer* (Girard).—Turtle Bay, April 4, two specimens about 100 mm long, seined. Off Abrejos Point, 17 to 20 fathoms, April 5, many caught on hook and line. Twenty-four specimens examined have 24 vertebrae each.

32. *Paralabrax clathratus* (Girard).—San Martin Island, April 1, 10 fathoms, one specimen on hook and line.

Xenichthyidae—Xenichthyid Fishes

33. *Xenistius californiensis* (Steindachner).—Magdalena Bay, April 7, two specimens 80 and 90 mm, seined.

Lutianidae—Snappers

34. *Lutianus guttatus* (Steindachner).—Between Seal Rock and Cerralbo Island, April 11, one specimen 625 mm long, hook and line. This example differs from descriptions in Meek and Hildebrand's *The Marine Fishes of Panama* in the smaller eye (5.75 in head); in the angulated anal fin (4th ray longest); and in the color, which in life was scarlet, slightly darker above axis; lower sides of head and body scarlet, with a silvery cast; a few light green spots anteriorly along axis of body below lateral line; belly and breast mostly white; fins plain scarlet; preopercle with violet cast; no large dark spot on back.

Haemulidae—Grunts

35. *Anisotremus interruptus* (Gill).—Turtle Bay, April 4, one specimen, 650 mm, seined.

Gerridae—Mojarras

36. *Eucinostomus californiensis* (Gill).—Turtle Bay, April 4, many seined. Magdalena Bay, April 7, many seined.

37. *Xystaema cinereum* (Walbaum).—La Paz, April 10, one specimen 330 mm, seined.

38. *Gerres lineatus* (Humboldt).—La Paz, April 10, one 220 mm, seined.

Kyphosidae—Rudder Fishes

39. *Girella nigricans* (Ayres).—Turtle Bay, April 4, four specimens seined, and many taken in tide pools. Socorro, April 9, twelve specimens from tide pools collected by Ira L. Wiggins. Halfway House, April 2, one specimen from tide pool, collected by Ira L. Wiggins.

Sciaenidae—Croakers

40. *Seriphus politus* (Ayres).—Turtle Bay, April 4, one specimen 145 mm, seined.

41. *Menticirrhus undulatus* (Girard).—Turtle Bay, April 4, two specimens 440 and 520 mm, seined.

42. *Cynoscion parvipinnis* Ayres.—La Paz, April 10, one specimen 520 mm long, seined. Two canines; eye 6.75 in head; pectoral 2 in head; dorsal X, I, 20. Seined. Apparently a new record for the Gulf.

43. *Genyonemus lineatus* (Ayres).—San Carlos Bay, April 2, twenty specimens, 200-300 mm, hook and line.

Embiotocidae—Surf Perches

44. *Hyperprosopon argenteum* Gibbons.—San Martin Island, April 1, one specimen 210 mm long, 10 fathoms, hook and line. San Carlos Bay, April 2, one specimen 170 mm, hook and line. A southern record.

Pomacentridae—Demoiselles

45. *Abudefduf marginatus* (Bloch).—La Ballena, April 30, three specimens from tide pools, collected by Ira L. Wiggins.

46. *Abudefduf declivifrons* (Gill).—La Ballena, April 30, one specimen from tide pool. Collected by Ira L. Wiggins.

Labridae—Wrasses

47. *Pimelometopon pulcher* (Ayres).—Off Abreojos Point, 17 to 20 fathoms, April 5, one specimen 400 mm long, hook and line.

Tetraodontidae—Swell Fishes

48. *Spheroides annulatus* (Jenyns).—Turtle Bay, April 4, one specimen 300 mm, seined. La Paz, April 10, twenty specimens about 80 mm in length, seined.

49. *Spheroides politus* (Girard).—La Paz, April 9, two specimens, 330 mm and 340 mm long, seined. These differ from specimens of *S. annulatus* in the much broader and flatter interorbital space (2.5 to 2.75 in head in *S. politus*, and 3.5 in *S. annulatus*) and in the coloration, *politus* having the black spots much more numerous and distinct.

Scorpaenidae—Scorpion Fishes

50. *Scorpaena guttata* Girard.—San Martin Island, April 1, two specimens, 210 mm and 290 mm long, hook and line.

51. *Sebastodes serranoides* Eigenmann and Eigenmann.—Off Cape Colnett, 43 fathoms, April 1, four specimens, hook and line. Apparently a southern record.

52. *Sebastodes goodei* Eigenmann and Eigenmann.—Ranger Bank, 70 to 90 fathoms, April 3, four specimens, hook and line. Thetis Bank, 60 to 70 fathoms, April 6, one specimen, hook and line. A southern record.

53. *Sebastodes pinniger* (Gill).—Off Cape Colnett, 43 fathoms, April 1, two specimens, hook and line. Apparently a southern record.

54. *Sebastodes elongatus* (Ayres).—Ranger Bank, 70-90 fathoms, April 3 and April 15, two and one specimens respectively, hook and line. Apparently a southern record.

55. *Sebastodes atrovirens* (Jordan and Gilbert).—San Martin Island, 10 fathoms, April 1, one specimen, hook and line. Apparently a southern record.

56. *Sebastes vexillaris* (Jordan and Gilbert).—Off Cape Colnett, 43 fathoms, April 1, one specimen, hook and line. Apparently a southern record.

57. *Sebastes chlorostictus* (Jordan and Gilbert).—Ranger Bank, 70 to 90 fathoms, April 3 and April 15, four and eight specimens respectively. Apparently a southern record.

58. *Sebastes constellatus* (Jordan and Gilbert).—Ranger Bank, 70 to 90 fathoms, April 3, three specimens, hook and line. Apparently a southern record.

Cottidae—Sculpins

59. *Clinocottus analis australis* Hubbs.—Turtle Bay, April 4, four specimens, tide pools.

Eleotridae—Sleepers

60. *Dormitator maculatus* (Bloch).—San Jose del Cabo, May 14, four specimens, collected by Ira L. Wiggins.

Gobiidae—Gobies

61. *Bathygobius soporator* Cuvier and Valenciennes.—La Paz, April 9, four specimens, 25-65 mm, tide pools.

62. *Gobiosoma histrio* Jordan.—Bahia de Concepcion, April 22, two specimens, collected by Ira L. Wiggins.

63. *Aboma chiquita* (Jordan and Evermann).—Bahia de Concepcion, April 22, nine specimens. Eight miles north of Santa Rosalia, April 21, three specimens.

Malacanthidae—Blanquillos

64. *Caulolatilus princeps* (Jenyns).—Off Cape Colnett, San Geronimo Island, Ranger Bank, Cerros Island, Abreojos Point, and Thetis Bank. April 2 to April 14. Abundant, hook and line.

Blenniidae—Blennies

65. *Labrisomus xanti* Gill.—La Ballena, April 30, one specimen, tide pool.

66. *Hypsoblennius gilberti* (Jordan).—Socorro, April 9, one specimen, tide pool.

67. *Runula asalea* Jordan and Bollman.—Magdalena Bay, April 6, one specimen, by drop-light. Mainland, opposite Cerralbo Island, April 11, three specimens, by drop-light, all about 35 mm long. Apparently the first record north of Cape San Lucas.

Ophidiidae—Cusk Eels

68. *Otophidium galeoides* (Gilbert).—La Paz, April 10, one specimen 75 mm long, on beach.

Pleuronectidae—Flounders

69. *Citharichthys fragilis* Gilbert.—Thetis Bank, 60 to 70 fathoms, April 6, one specimen 220 mm long, hook and line. Apparently the first record from the west side of the peninsula.

NATURAL HISTORY MUSEUM, STANFORD UNIVERSITY, CALIFORNIA.

Feeding Habits and Molt of *Crotalus confluentus oregonus* in Captivity

By TRACY I. STORER and BERYL M. WILSON¹

THERE is a notable dearth of exact information concerning the food habits of serpents. Data on these reptiles in the wild consist chiefly of isolated observations on prey actually observed in process of being captured or eaten, or of the food found in the stomachs of specimens which have been killed. The slight chance for observations of the first sort and the large number of specimens examined in which the digestive tract is empty have further limited the number of useful records. The chance receipt of three living specimens of the Pacific rattlesnake (*Crotalus confluentus oregonus* Holbrook) in a laboratory where a colony of domestic Norway rats is maintained for nutritional studies has afforded opportunity to study the food habits of this species.

Our first specimen (No. 1) was captured at Bluff Lake, 7,500 feet altitude, San Bernardino Mountains, California, on June 26, 1928. This animal was in the "black phase" of coloration. The other two were received May 6 (No. 2) and in June (No. 3) respectively, of 1930; one of these (No. 2) was obtained somewhere in the San Joaquin Valley; the other presumably somewhere to the north and west of Davis: it was impossible to learn the exact localities of capture. However, all are undoubtedly of the sub-species indicated above.

No. 1 weighed 452 grams on August 25, 1928. No measurements have been taken, and, unfortunately, exact record of the rattles has not been kept. No. 1 had either 6 or 7 rattles when received; it has lost some, at the tip, and now has 10 rattles. It has molted twice each year or 7 times in $3\frac{1}{2}$ years and so should have either 13 or 14 rattles if one is acquired at each molt.

However, close inspection of the snakes after certain molts has raised a suspicion in our minds that they did not always acquire a rattle with each molt; this needs confirmation. Snake No. 2 molted twice in 1930 and once in 1931, while No. 3 molted once in 1930 and twice in 1931. All of the snakes have lost terminal rattles, but in the absence of distinguishing marks on the individual rattles the number so lost is uncertain.

The eyes of snake No. 1 appeared "milky" or "clouded" on August 3, 1931; this condition had cleared up by August 6 and the snake molted on August 12. With No. 2 the eyes appeared clouded on June 13, 1931, and the snake molted on June 23. In three years the first molt of the year of snake No. 1 has occurred from May 27 to June 8, and the second molt (in 4 years) from July 27 to August 20. Here then is a suggestion as to the basis for the popular belief that "rattlesnakes are blind in August," that is, preparing to molt. The molts of the other two snakes have been somewhat more irregularly spaced.

¹Contribution from the Division of Zoology, College of Agriculture, University of California.

There has always been a decided difference in the temperament of these three snakes. No. 1 has from the beginning of its period of captivity been "sullen" and has rattled on the least provocation. When first in the laboratory it rattled almost continuously. At that time the snake was caged in a room where the step of a person vibrated the floor and the snake responded by rattling at any approach. Then, also, a low vocal note from a person would stimulate it to rattle. Now, in a more quiet and substantial location, and after more than $3\frac{1}{2}$ years in captivity, it still responds to a slight disturbance by rattling. This snake spends much of its time in the coiled position, and if, when approached, it happens to be outstretched, it immediately assumes the coiled position and commences to rattle. The other two, paler, brown and yellow colored, snakes have reacted differently. They seem to require much more stimulus before beginning to rattle. No. 2 (the larger) spends much of its time extended and seems to exhibit curiosity when approached, coming up to the glass front of the cage when an observer places a hand on the outside of the glass; this snake always follows any cleaning implement introduced into the cage. When moving about in its cage (11" high by 24"x36" floor), it often travels with its head close to the top of the enclosure, whereas the other two, when moving, keep their heads and bodies close to the shavings on the floor of their respective cages.

Our observations on the difference in reaction of the blackish snake (No. 1) are paralleled by the experience of a highway-construction foreman on the Big Trees road east of Dorrington, Calaveras County, California, who stated that in the summer of 1930 his crew had killed between 25 and 30 rattlesnakes (all *oreganus*) in that region; all were relatively docile except one which he characterized as a "black diamond rattler" and that individual showed much fight before being dispatched.

Our No. 1 snake when first captured was almost black. The dorsal pattern was relatively inconspicuous. Prior to its earlier molts it assumed a slightly more grayish appearance, but when fresh molted was almost velvety black again. Now, within the past year, it has become much grayer and has maintained this tone of coloration for a long period of time.

Since their respective arrival in the laboratory all of these rattlesnakes have been offered food in the form of live rats at frequent intervals. We found by experience that if offered food while an observer was present the snake would usually kill the rat but would not eat it. In consequence it has been our practice to place rats in the cages at the end of the day, so that the snakes would be undisturbed when ready to feed. Water was offered snake No. 1 in 1928, but there was no evidence that any was taken so none has been provided for any of the snakes subsequently. Other reptiles in captivity are known to require drinking water. The water necessary for these rattlesnakes has, therefore, been obtained entirely from the fluids in the bodies of the rats consumed. The dry substance in the body of the captive Norway rat averages around 32 per cent, hence the fluid (water) is about 68 per cent (Donaldson, H. H., *The Rat*, revised ed., 1924: 300). The total available fluid intake for snake No. 1 has, there-

fore, been 705, 983 and 980 grams, in 1929, 1930, and 1931 respectively, or a total fluid intake per year but slightly in excess of the body weight of the animal itself. A human being if restricted to the same proportion of fluid intake would receive about twenty gallons annually! Fluids are lost from the body of a reptile in small amounts in the feces, in the secretions involved in sloughing of the skin, and in the respiratory process. Presumably there is no water loss through the skin. The body temperature of a reptile is unregulated; it is in most instances practically that of the environment. Lacking a regulated body temperature there is a lessened need for moisture, which, in birds and mammals, is used to a considerable extent in the regulation of body temperature. The success of this degree of conservatism in water maintenance of the rattlesnake is further indicated in an instance reported by Duméril and quoted by Flower (*Proc. Zool. Soc. London*, 1925: 977), of a specimen in the Jardin des Plantes in Paris that went 22 months without eating and then fed and continued to live for 10 years. The water relations of our snakes at Davis are all the more remarkable since these snakes have been kept in a room moderately heated in winter and exposed to the heat of mid-summer in the Sacramento Valley, tempered only by windows and the walls of a reinforced concrete building. It should be noted that Mitchell (*Researches upon the Venom of the Rattlesnake*, Smiths. Contr. to Knowledge, 1861: 4) reported the necessity of a water supply for his snakes, especially in respect to molt. We have not had this experience. Our snakes, with the possible exception of one instance with No. 1, have molted their skins entire. Mitchell was using *Crotalus durissus* [= *horridus*], an inhabitant of the eastern humid region, while we have used *oreganus*, an inhabitant of the semi-arid region of California.

Of rats offered and not killed by the snakes, several have subsequently been killed by us, but we have found no evidence that they had been struck save in one instance. One rat offered April 19, 1929, to snake No. 1, was struck in the middle of the back. This rat was still alive on the following day and when killed by us and examined showed only two fang punctures in the skin, surrounded by darkened areas; the muscles adjacent and the viscera appeared normal. The snake had killed a 91 gram rat at 5 P.M. on April 11, which was eaten after 12 M. on April 12.

These snakes are all kept in solid wooden cages, the floors of which are covered with wood shavings, changed at long intervals. Record has been kept of every fecal mass, as noted in the chart. It is evident that feces are not passed after each feeding, but only at long and irregular intervals.


Detailed record of the food, feces and molt has been kept since these snakes arrived in the laboratory. In summary, snake No. 1 in 1928 ate, between July 20 and October 3, four rats and two house mice, totalling probably 400 grams (records not complete in 1928), voided feces twice and molted once; in 1929 it consumed 11 rats, totalling 1,037 grams in weight during 266 days, voided feces four times and molted twice; in 1930 it ate 17 rats totalling 1,446 grams, in 264 days, voided feces once and molted

twice; and in 1931 consumed 7 rats grossing 1,442 grams, in 170 days, voided feces 4 times and molted twice.

Snake No. 2 in 1930 ate 15 rats (all under 100 grams in weight) totalling not over 952 grams, between May 15 and August 12 (89 days), and shed twice; in 1931 it ate 5 rats, weighing 861 grams, in 71 days, voided feces 4 times and molted once.

Snake No. 3 in 1930 ate 9 or 10 rats, grossing between about 550 and 600 grams, in 66 days and molted once; in 1931 it ate 5 rats totalling 768 grams, in 99 days, voided feces twice and molted twice.

The exact amounts taken by snakes Nos. 2 and 3 in 1930 are slightly in doubt as these two snakes are housed in one cage and it was not possible in some instances to tell which of certain small rats put in the cages were eaten by each of the snakes. In our notes there is no record of feces passed in 1930 by snakes Nos. 2 and 3, but there is at least a possibility that these were not recorded by the attendant.



Record of food taken (L), or rejected (T), feces passed (O) and molt (X) of three captive specimens of *Crotalus confluentus oregonus* Holbrook

The size of rats offered for feeding has varied. In 1929 No. 1 ate rats of from 67 to 128 grams weight; a few of larger weight were not eaten but this may not have been due to size so much as satiety. In 1930 the weight of rats fed varied from 52 to 155 grams, while in 1931 distinctly heavier rats were taken, of 104 to 326 grams; others refused in this year were within these limits. The other two snakes have eaten smaller rats, below 100 grams in 1930 and from 96 to 204 grams in 1931. It is not yet evident whether some of the larger rats offered these two snakes in 1931 were refused because of larger size or because the snakes had ceased to feed for the season, although we are inclined to the latter explanation since many of those offered but not eaten were well within the weight range of those taken earlier in the year.

These data on the Pacific rattlesnake in captivity obviously do not give an accurate idea of the amount of food required by the species in the wild; yet they do indicate that relatively small amounts of food with fluid

content as calculated, are adequate not only for maintenance, but provide for growth and molt. Even if doubled or trebled under natural conditions the totals would be very modest in relation to the necessary food intake of homiothermous birds and mammals, and would explain why so many of the snakes collected in natural surroundings have no recognizable food materials in the digestive tract, and likewise the apparent scarcity of serpents in many places. Taking food in relatively large units only at considerable intervals they may remain in retirement and inactivity much of the time, a condition decidedly in contrast to that of birds and mammals much of whose time must of necessity be spent in procuring food.

The accompanying chart gives a graphic record of activities of these snakes in captivity.

UNIVERSITY OF CALIFORNIA, DAVIS, CALIFORNIA.

Blanding's Turtle, *Emys blandingii* (Holbrook), in Pennsylvania

By M. GRAHAM NETTING

WHEN as well-informed a worker as De Sola¹ omits Blanding's turtle from consideration in a list of the turtles of the northeastern states, as noted by Babcock,² it seems worthwhile to call attention to the occurrence of this species in Pennsylvania. The first record of *Emys blandingii* in this state, based on specimens in hand, was published by Stewart.³ His record was based upon two specimens which were collected near Lewisburg. In correspondence, Dr. Stewart informed me that one of these specimens was brought to him in April or March of 1927 from Kincade Swamp, which lies in Northumberland County although only one mile east of Lewisburg. The second specimen consisted of a carapace only which Dr. Stewart found in the collection of the Bucknell University Museum, and which one of his associates recalled as having been taken about 1905 in a small pond one mile south of Winfield in Union County. The distance between these localities is approximately five miles. In addition to the above-mentioned specimens known from Pennsylvania, the Carnegie Museum has two others; No. 3188 was collected at Conneaut Lake in Crawford Co., on August 20, 1904, by Dr. D. A. Atkinson, and No. 3189 was collected at Linesville, in the same county, on June 9, 1906, also by Dr. Atkinson. These localities are approximately six miles apart.

Numerous hypotheses, none of which can be accepted as final, may be offered in explanation of this discontinuous distribution.

1. Blanding's turtle may be found elsewhere in the northern part of the state if systematically hunted over a period of years.

¹ De Sola, C. Ralph; Bull. N. Y. Zool. Soc., 34, (5), 1931: 131-159.

² Babcock, Harold L.; Copeia, (1), 1932: 43.

³ Stewart, N. H.; Proc. Penna. Acad. Sci., II, 1928: 24.

2. The records for the Lewisburg area may indicate that stream capture at the western headwaters of the Susquehanna permitted this turtle to cross from the Allegheny system together with *Cryptobranchus alleganiensis* and *Graptomys geographica*.

3. The Lewisburg specimens may be the result of human introduction. In a letter to me Dr. Stewart writes, "It is possible that in some years past teachers of zoology here may have liberated some imported material. I can realize that that might have happened in the case of the one from Kincade Swamp, though I have not released any in twenty years. I doubt very much if the Union County specimen had ever been in captivity. I hope to find some in places that would not admit of any possibility of importation." The fact that Blanding's turtle was used widely in the past in comparative anatomy classes lends credence to this view. Since both of the records are from points close to the Susquehanna River, marshy spots that probably were at one time lagoons of this river, it is quite possible that a few escapes established themselves in this region and spread along the river to suitable localities.

4. The Carnegie Museum specimens from the Pymatuning region may indicate that the species is a relict form which persists in this boreal refuge. Many species of plants and animals, of which *Clemmys muhlenbergii* is one, exist as relicts in the Pymatuning area although rare, absent, or spottily-distributed elsewhere in the state. Most herpetologists may be expected to favor this explanation.

5. I must confess, however, that I find myself intrigued by my final hypothesis; namely, that Blanding's turtle may have reached Crawford County by way of the Beaver and Lake Erie Canal. This canal was completed in 1844 and closed in 1871, with an aftermath of rigorous destruction, so that its life as a continuous water-course can scarcely have exceeded thirty years. Yet in this short period *Emys blandingii* may have traversed the forty-five miles of water from the shore of Lake Erie, where it still occurs, to Conneaut Lake, which was directly connected with the canal. The Linesville specimen was taken at a locality about four miles west of the old canal bed.

I offer this suggestion that a canal may have influenced the local distribution of a turtle, not because it can be proved at the present time, but because it may stimulate zoögeographers in contiguous states to examine their maps in the light of the possible effect, on semi-aquatic or aquatic biota, of the network of canals which existed during the middle nineteenth century. In those days of canalization four major water-ways had outlets on Lake Erie: one extended southward from Toledo; another extended southward from Cleveland; the third connected Erie and Beaver; and the fourth, and most famed, extended from Buffalo eastwards. Certain of the canals were joined by short "feeders" or "crosscut" canals. Most of them were in existence for extremely few decades, yet we know that the Panama Canal during its short period of existence has begun to act as a pathway for the migration of certain fish.

CARNEGIE MUSEUM, PITTSBURGH, PENNSYLVANIA.

Herpetological Records and Notes from the Vicinity of Tucson, Arizona, July and August, 1930

By F. WILLIS KING

AMPHIBIANS

1. *Scaphiopus couchii* Baird

This spadefoot was found in greatest numbers after heavy rains in puddles and backwater along the Rillito River near University Farm. The height of their breeding season was between July 11 and 15. Their call, which resembled most a plaintive bellow, began shortly after dark and continued until just before dawn. They call both from the edge of the muddy water and while swimming, the latter call being muffled: the male exerts his whole body in producing his call, drawing in his hind legs as the vocal pouch is protruded. The eggs varied from six to twenty-four in a clump, and are fastened in irregular masses to any solid body in the water, just below the surface, hatching in one or two days. The little tadpoles grow rapidly and are sufficiently developed in a week to make their way into the mud at the bottom of the puddle and all have vanished by two weeks.

The mucus from the spadefoot is irritating to any break in the human skin, and has a pungent odor. The males outnumber the females four to one, are more active, and are often found quite a distance from water. By August 15 the species had disappeared.

2. *Scaphiopus hammondi* Baird

This species was almost always found in association with the preceding one, but in fewer numbers. The breeding season was closely correlated with that of *S. couchii*. While the adults range from 50 to 60 mm. in body length, one young female measuring only 33 mm. was taken which showed well developed eggs. The call is more vibrant than that of *S. couchii*. This spadefoot is an active, trim, creature.

3. *Bufo alvarius* Girard

This large toad was common around the barns and gardens at the University Farm. No call was heard or breeding activities observed during July and August. The species is of considerable economic importance because of the great quantity of June and fruit beetles it catches, as well as spiders and grasshoppers. It also feeds on small lizards. I have handled numbers of them, and have seen others do likewise with no ill effects. The body color varies from a bluish gray to green or brown. The smaller warts of the back are often of a rusty color, as are the tips of the digits. The belly is white. The animals are often found in open pits, holes or washouts where they have fallen. One specimen I have just fills a quart jar.

4. *Bufo cognatus cognatus* (Say)

This is the most common toad of the southwest. These toads call and breed in and along the irrigation ditches, whenever water is present. They

were seen in great numbers during July. They leave the ditches and wander through the fields and gardens between rains and irrigation intervals and show little fear of man. As is usual among toads, the males are smaller but more numerous than the females. The digging spur is well developed.

5. *Hyla arenicolor* Cope

Tree toads of this species were found in White House Canyon, in the Santa Rita Mountains, August 3. Males were heard calling just before dark from branches of oak trees, about fifteen feet from the ground. The call was an even, strong trill. Tadpoles of *H. arenicolor* were found in pools nearby with some of the larvae transforming at 18 mm. body length.

6. *Rana pipiens* Schreber

Leopard frogs were common along the stream in Sabino Canyon. One large individual was also found under a bunch of hay on University Farm, at least two hundred yards from any permanent water. The young were transforming the latter part of July. All individuals examined showed very bright orange coloration on backs of thighs and on flanks. The dorsal spotting is irregular; the tympanum may or may not have a light spot; the spot on the snout is also variable; the legs are barred. The general coloration is grayish with many showing green or brown as a dominant color.

REPTILES

1. *Cnemidophorus sexlineatus perplexus* (Baird and Girard)

Common in Santa Catalina foothills. Two forms of this species are found. One shows spotting between the longitudinal stripes and on the thighs. The distance between the median stripes equals the distance between the first and second stripes on the side. The other form has the spotting indistinct or absent, and the median dorsal stripes very close together. These variations are not correlated with size.

2. *Uta stansburiana elegans* (Yarrow)

Taken on University Farm.

3. *Uta ornata linearis* Baird

Common on University Farm.

4. *Holbrookia maculata approximans* (Baird)

The specimen of *Bufo alvarius* referred to above regurgitated my only specimen of this lizard.

5. *Holbrookia texana* (Troschel)

Common in the Santa Catalina foothills in association with *Callisaurus ventralis*.

6. *Callisaurus ventralis ventralis* (Hallowell)

A conspicuous lizard of the Santa Catalina foothills.

7. *Sceloporus clarkii* Baird and Girard

One specimen was taken from a stone wall in White House Canyon, Santa Rita Mountains, August 2.

8. *Sceloporus jarrovi* Cope

Common in the rocky areas of White House Canyon between 6,000 and 8,000 feet elevation. Young individuals are darker than the adults. Many are infested with red mites.

9. *Phrynosoma solare* Gray

Common in the latter part of July along Prince Road, north of Tucson.

10. *Phrynosoma douglasii hernandesi* (Girard)

One individual was taken at an elevation of 9,000 feet on Mount Baldy in the Santa Ritas, August 3. Its colors harmonized perfectly with the reddish brown and gray of the soil and rocks.

11. *Heloderma suspectum* Cope

The Gila monster is still quite common in the rocky foothills of the ranges surrounding Tucson. Occasionally one is seen awkwardly crossing a road or is found in a garden. I found them in the foothills of the Santa Ritas and Catalinas.

12. *Chilomeniscus cinctus* Cope

One specimen was found under a stump near Prince Road, August 15. This snake could move through loose dirt as fast as it could over the surface.

13. *Tantilla nigriceps* Kennicott

One of these small snakes was taken from a swimming pool in White House Canyon, August 3.

14. *Masticophis flagellum frenatus* (Stejneger)

A large individual was observed in White House Canyon, August 3.

15. *Coluber constrictor mormon* (Baird and Girard)

Found along irrigation ditches, University Farm.

16. *Pituophis catenifer rutilus* Van Denburgh

An individual four feet in length was found on North Campbell Avenue.

17. *Terrapene ornata* (Agassiz)

This box turtle was common in the gardens on University Farm. An examination of stomach contents showed its food to be largely insects, chiefly June beetles.

REFERENCE LIST

BLANCHARD, FRANK N.

1925 A Key to the Snakes of the United States, Canada, and Lower California. *Pap. Mich. Acad. Arts, Sci. and Lett.*, 4, Part 2: 81 pp.

DICKERSON, MARY C.

1906 The Frog Book. Doubleday, Page and Co.: 280 pp.

SLEVIN, JOSEPH R.

1928 The Amphibians of Western North America. *Occ. Pap. Calif. Acad. Sci.*, 16: 152 pp.

VAN DENBURGH, JOHN

1922 Reptiles of Western North America, 2 vols. *Ibid.*, 10:1028 pp.

BIOLOGY DEPARTMENT, WILMINGTON COLLEGE, WILMINGTON, OHIO.

Herpetological Notes

THE STATUS OF TWO PERUVIAN LIZARDS.—The discovery that one of the specimens determined by Boettger (1889, p. 311) as *Amphisbaena occidentalis* Cope was conspecific with examples collected close to the type-locality of *A. townsendi* Stejneger, raised doubts as to the correctness of Boettger's determination or the validity of the last-mentioned species. In the description of *A. townsendi* a number of differences from *occidentalis* are mentioned, but only one of these appears to be significant,—the possession by the former of about 100 more annuli on the body. For many years this alleged disparity could not be checked owing to the disappearance of Cope's types, but now, thanks to the researches of Professor Dunn they have been re-discovered in the museum of the Philadelphia Academy, and the counts, which he has courteously made at the writer's request, reveal that Dr. Stejneger was the victim of a miscount. *A. townsendi* is probably conspecific with *A. occidentalis*, but the very limited series of specimens available makes it seem possible that a northern race with a slightly higher number of caudal annuli may prove to be recognizable.

Specimens	Locality	Annuli on Body	Annuli on Tail
Cotypes of <i>A. occidentalis</i>	Valley of Jequetepeque	276	19
Cope		273	23
Phil. Ac. 11355-11358		266	21
		269	21
Boettger's specimens	Pacasmayo. "Only a few	273	18
(Boettger 1889) B.M.N.H. 89.	Kilometers from Cope's	265	22
7.19.1.	original locality."	278	23
		274	25
<i>A. townsendi</i> Type. U.S.N.M.	Piura	276	25
47087.			
		277 (approx.)	25
B.M.N.H. 1929. 12.12.2-5	Lobitos, Talara	271	26
coll. Dr. Burgess Barnett.		277	27
		277 (approx.)	28

The specimens called *A. darwini* by Peracca (1895, p. 9), but thought by Stejneger to be referable to *A. occidentalis*, have already been referred to a distinct species (Parker 1928, p. 383).

REFERENCES

- BOETTGER, O.
1889 Ber. Senck. Nat.forsch. Ges.: 311.
COPE, E. D.
1876 Journ. Acad. Sci. Philad., (2) 8: 176.
PARKER, H. W.
1928 Ann. Mag. Nat. Hist., (10) 2: 383.
PERACCA, G. M.
1895 Boll. Mus. Torino, 10, 195: 9.
STEJNEGER, L.
Proc. U. S. Nat. Mus., 41: 283-4.

H. W. PARKER, *British Museum (Natural History), London, England.*

COMMENTS ON POLYPEDATIDAE.—I have recently had the opportunity of examining a considerable number of specimens of those genera of frogs which Noble has recently grouped into a new family, Polypedatidae. These investigations have been carried on in the extensive collection at the Museum of Comparative Zoology.

The only consistent character differentiating this group of genera from those of the Ranidae is the presence of an intercalary cartilage in the digits and an examination of several species in each of eleven of the thirteen genera included in this group reveals the presence of intercalary cartilages in every case. Although not all of the species available were examined, and although two genera were omitted, the results are nevertheless significant.

The following list of species were examined:

Aglyptocephalus madagascariensis (A. Duméril);

Chiromantis petersii Boulenger, *rufescens* (Günther), *xerampelina* Peters;
Hylambates hyloides Boulenger, *maculatus* A. Duméril, *natalensis* (Smith);
Hyperolius concolor (Hallowell), *horstocki horstocki* (Schlegel), *kivuensis* Ahl,
marginatus Peters, *mariae* Barbour and Loveridge, *narmoratus* Rapp, *montana*
Angel, *ocellatus* Günther, *platyrhinus picturatus* Peters, *platyrhinus* Procter,
puncticulatus (Pfeffer), *rhodoscelis* (Boulenger), *rossii* Calabresi;
Kassina senegalensis (Duméril and Bibron);
Leptopelis aubryi (A. Duméril), *bocagii* (Günther), *brevirostris* Werner, *johnstoni*
(Boulenger), *notatus* (Bucholz and Peters), *ocellatus* Mocquard, *palmatus*
(Peters), *rufus* Reichenow, *uluguruensis* Barbour and Loveridge;
Mantella baroni Boulenger, *ebenau* (Boettger);
Mantidactylus guttulatus Boulenger, *ulcerosus* (Boettger);
Megalixalus brachynemis Boulenger, *dorsalis* (Peters), *forasini* (Bianconi), *madagascariensis* (Günther);
Philautus aurifasciatus (Schlegel), *glandulosus* (Jerdon), *leucorhinus* (Martens),
vittatus Boulenger;
Polypedates aculeatus Gray, *appendiculatus* Günther, *bimaculatus* Boulenger, *eques*
Günther, *goudoti* (Tschudi), *leucomystax* (Gravenhorst), *macrotis* Boulenger,
maculatus (Gray), *mocquardi* Boulenger, *ostoni* Stejneger, *pardalis* (Günther),
reinwardti (Wagler), *schlegeli* Günther, *tephracomystax* A. Duméril.

It is worthy of notice that some of these, such as *Mantidactylus guttulatus*, are large frogs, others, like *Megalixalus brachynemis*, are small. Examination of a number of Raninae, Petropedatinae, and Cornuferinae failed to disclose any intercalary cartilages.

The presence of an intercalary cartilage is a consistent taxonomic character for this group. It is not, however, in my opinion really worthy of elevation to the status of a family but rather it seems to me to be characteristic of a very good subfamily of the Ranidae. The presence or absence of intercalary cartilages is a character more comparable with such characters as the presence or absence of vomerine teeth, of calcification characters of sternum and omosternum, the shape of the pupil, the presence or absence of digital dermal scutes, and the shape of the terminal phalanges rather than with more fundamental characteristics of skeletal variations on which family differences had best be considered to be dependent.

Noble claims that this probably natural group has evolved from the Ranidae in much the same way that the Hylidae have evolved from the Bufonidae. It is true that a case of parallel evolution appears to exist between these two widely separate groups but would not there be numberless families in the classification of amphibia if they were based on instances of parallel evolution?—BENJAMIN B. LEAVITT, *Museum of Comparative Zoology, Cambridge, Massachusetts.*

AN ABERRANT *NATRIX CLARKII* (BAIRD AND GIRARD).—A female specimen of *Natrix clarkii* (M.J.A., No. 1304), taken on the back bay of Biloxi, March 18, 1932, from a typical salt marsh habitat, exhibits certain peculiarities in color pattern that, while of probably no particular significance, are considered of sufficient interest to warrant recording, especially in view of the fact that with the exception of this individual all other specimens coming under my observation have been constant in their agreement with the characters of the species.

In other respects the specimen appears to be normal in having 21 scale rows, 10 infralabials, 8 supralabials and 134 gastrosteges. It is 367 mm. in length. The coloration is much faded but the pattern consists of a faint median dorsal line extending from the parietals to 5 scale rows posterior to the anus and covering the middle row of scales and half of each adjoining scale row. The lateral stripes are more distinct and anteriorly are 1 to 1½ scale rows in width but 115 mm. from the snout become broken and alternately expand and contract until at a point 75 mm. in front of the vent they become joined to the light band, encompassing the tips of the gastrosteges and first row of scales by a series of hour glass-shaped markings of a brownish hue. These are 1 to 2 scales wide at the constriction and broaden to 2 to 4 scales in width at their junction with the longitudinal lines. The lowermost stripe is similar to the second in its irregularity.—MORROW J. ALLEN, *Biloxi, Mississippi.*

Ichthyological Notes

FISH NOTES FOR 1931 AND 1932 FROM SANDY HOOK BAY.—The work of the New York Aquarium collecting boat, *Seahorse*, brought to light no new records or items of special interest in 1931. Consequently, all the data here listed refer to the year 1932.

1. *Hyporhamphus roberti*. (Cuvier and Valenciennes).—A new record for Sandy Hook Bay. Two specimens, both males, 144 and 150 mm. in standard length. These were taken August 11, in company with a school of *Strongylura marina* (Walbaum) of about the same size, and were seined together with *Synodus foetens* (Linnaeus), *Mullus auratus* Jordan and Gilbert and *Synynathus fuscus* Storer.

2. *Vomer setapinnis* (Mitchill).—Small specimens of this species are generally common in Sandy Hook Bay. These range up to about 50 mm. in standard length. In 1932, during August and September, however, much larger ones were also common. A sample of the larger fishes (17 specimens) ranged from 141 to 167 mm. in standard length and averaged 152 mm.

3. *Trichiurus lepturus* Linnaeus.—A single specimen of 876 mm. total length was taken in a pound net on June 30. It is a new record for this Bay, as well as both an early and a size record for the New York region.

4. *Chaetodon ocellatus* (Bloch).—A single specimen, 34 mm. in standard length; October 26. A late date.

5. *Mola mola* (Linnaeus).—A single specimen with the following measurements was taken on September 15. Total length, 126 cm.; total height, 160 cm., weight, 227 pounds. It was an unripe male. The stomach contents consisted of three jellyfish, *Aurelia* sp., various seaweeds including *Ulva*, and some brown algae, as well as fragments of sponge. This specimen was turned over to the American Museum of Natural History. A late date for Sandy Hook Bay.

The unusual run of small mackerel about New York in 1932, did not seem to get into Sandy Hook Bay in any quantity although they were taken in numbers from State Island piers and elsewhere, as far east as Montauk Point. They were most abundant about New York City in the first week of August. Prior to that date, July 12, still smaller ones were abundant at the Montauk Yacht Club harbor¹. On that date a sample of 12 *Scomber scombrus* Linnaeus ranged from 127 to 143 mm., and averaged 138 mm. These were apparently some of the same group that later appeared in New York Harbor.

This list adds two species to the fish fauna recorded from Sandy Hook Bay, *Hyporhamphus roberti* and *Trichiurus lepturus*, bringing the total to 122. The series of records reported mostly in COPEIA for the last twelve years, if divided into quarters, shows the following numerical relationship with regard to the total number of species recorded at the end of each period: 95, 112, 118 and 122. From this it would seem that slight additions can be subsequently expected.—C. M. BREDER, JR., *New York Aquarium, Battery Park, New York City*.

THE STATUS OF *OSPHRONEMUS SAIGONENSIS* BORODIN.—In the Bulletin of the Vanderbilt Marine Museum, vol. I, art. 2, 1930, p. 48, Borodin has described this species, based on a single specimen collected at Saigon, French Indo-China. It is evident from the description that the fish is not an *Osphronemus* but a *Trichogaster*. As I am very familiar with the four recognized species of this genus, all of them abounding in Siam, I am interested in the reported existence of another species in Indo-China, where at least three of the known forms occur. Borodin notes that the species nearest to *Osphronemus saigonensis* is *O. siamensis* Günther, which is a synonym of *Trichogaster trichopterus* (Pallas). This is a very common fish in Siam and in Cambodia and Cochin China (of which last Saigon is the capital), and is subject to considerable variation in shape, squamation, fin formulae, and color. This variation would apparently fully cover *saigonensis*.—HUGH M. SMITH, *Department of Fisheries, Bangkok, Siam*.

¹These notes were made on a cruise of the *Querida*, while guest of Mr. Daniel Bacon.

A NEW RECORD OF *OSTRACION DIAPHANUM* BLOCH AND SCHNEIDER FROM CALIFORNIA.—Three specimens of the cowfish or trunkfish, *Ostracion diaphanum* Bloch and Schneider, have recently been taken along the coast of Southern California and brought to the Los Angeles Museum. This appears to be the first occurrence of the species on the Pacific Coast of the United States. Its usual range is from the East Indies to Japan. The present examples are typical in form and coloration with the exception that some variation is shown in the structure and arrangement of certain spines on the carapace.

The largest specimen was taken alive in a net off Santa Barbara, California, the latter part of November, 1932. It measured 7 inches in length, the carapace being $4\frac{3}{4}$ inches long. The left supraocular spine is bent more sharply outward than is the right. The lateral spine on the right dorsal ridge is larger and more advanced forward than the corresponding spine on the left dorsal ridge. The median dorsal spine is scarcely recurved.

Another individual was caught in a net near the kelp beds off San Pedro, California, on December 1, 1932. Its total length is 5 inches and the carapace $3\frac{1}{4}$ inches.

The other specimen was washed up on the beach at Santa Monica, California, on December 5, 1932. It measures $5\frac{1}{4}$ inches in length and the carapace $3\frac{3}{4}$ inches. The left supraocular spine is undeveloped while the corresponding spine on the right is short and much thickened. The central, lateral spine on the right ventral ridge is weakly developed and is only represented by a small, sharp point. On the left ventral ridge, the position of the central spine is nearer to the posterior spine than to the anterior. The normal position of the central spine is half way between the two.

In all specimens, several rows of black blotches, roughly circular in outline, appear above and below the ventral ridge. Otherwise, the lower part of the body, below a line drawn from the gill opening to the tail, exhibits the characteristic translucent appearance of the species.—HOWARD R. HILL, *Los Angeles Museum, Exposition Park, Los Angeles, California.*

A RECORD OF THE LUVARU (*LUVARUS IMPERIALIS* RAFINESQUE) FROM SOUTHERN CALIFORNIA.—A large specimen of the rare *Luvarus imperialis* was captured at Redondo Beach, California, near Los Angeles, on the afternoon of November 20, 1932. It measured six feet, one inch, in length and weighed 305 pounds. Only once before has this species been recorded from local waters. The fish came ashore, in a live but apparently injured condition and was killed with an iron bar. It was taken to the Los Angeles Museum where a life size cast was made of the body and the skeleton prepared for exhibition.—HOWARD R. HILL, *Los Angeles Museum, Los Angeles, California.*

A CASE OF ARTERIAL CONSTRICTION OF THE SPLEEN IN THE DOGFISH.—Recently a large male of the dogfish shark (*Squalus acanthias*), which was under dissection in our comparative anatomy laboratory, was found to have the spleen divided into a small posterior lobe and a large anterior lobe by a transverse constriction formed by three loops of the posterior intestinal artery. While there is apparently no way of determining whether this abnormality was developed in embryonic life or not, it had probably been present for a considerable time, since the two splenic lobes were held together by a narrow cord-like constriction less than two millimeters wide. The greatest total length of the organ was 103 mm., the length of the anterior lobe being 68 mm., and that of the posterior lobe 35 mm. The greatest width of the anterior piece was 35 mm., as contrasted with 12 mm. for the posterior section. No differences in color were noted when the two lobes were compared. Numerous small blood vessels were visible on the surface of the connecting cord, in which they were concentrated due to the pressure of the arterial strangulation. The three loops in the posterior intestinal artery were easily unwound and the spleen freed of this feature during the present examination of the specimen, at which time it was apparent that due to the natural slack in this artery these coils had not been pulled taut enough to accomplish complete amputation of the tip of the spleen.—CHARLES E. BURR, *Southwestern College, Winfield, Kansas.*

BLUE PERCH.—On September 3, 1932, Mr. W. D. Bates of Ridgetown, Ontario, sent to the Royal Ontario Museum of Zoology a specimen of what he called albino perch. The specimen was packed in ice, and was received in excellent condition. On receipt it appeared bluish, suggesting the color of the blue pickerel, *Stizostedion glaucum* Hubbs. In a subsequent statement Mr. Bates said that the perch had the bluish shade when caught, that it was marked by the black, vertical markings as in the case of the typical form, but that there was no yellow coloration in it. He also said that perch of this coloration are very rare, and that he had probably seen only half a dozen in his fifty years' fishing in lake Erie.

One cannot help thinking that the cause of this bluish coloration in a typically yellow colored species is similar to that which gives rise to the blue pickerel. This difference is probably in the nature of a physiological variation, and I believe that specific differentiation often has its inception in such variations. Whether natural selection will ultimately produce a blue perch is something which only time can tell.

Dr. Carl L. Hubbs informs me that he has had a trustworthy report of similar blue perch occurring rarely in Saginaw Bay, Michigan.—J. R. DYMOND, *Royal Ontario Museum of Zoology, Toronto, Canada.*

REVIEWS AND COMMENTS

PROGRESS ON THE BASHFORD DEAN MEMORIAL VOLUME ON ARCHAIC FISHES.—Soon after the death of Professor Bashford Dean on December 6, 1928, some of his former colleagues and friends at the American Museum of Natural History and Columbia University met and formed the Bashford Dean Memorial Committee. The object of the committee was to bring about the publication of a memorial volume containing among other appropriate matters a series of color plates reproduced by photolithography. The original drawings for such plates had been executed chiefly by Dean himself for his own projected memoirs on the embryology of the California hagfish (*Bdellostoma stouti*), the Frilled Shark (*Chlamydoselachus anguineus*) and the Port Jackson Shark (*Cestracion* (= *Heterodontus philippi*). This exceedingly rare and important series of eggs and embryos had been collected by Dean in several parts of the world during his long activities as a zoologist. A few of them he had already published, but he had been prevented from completing and publishing the greater part of his material by the circumstance that he was equally eminent and probably much more widely known as a collector and connoisseur in European armor of the middle ages and that he finally became the founder and curator of the Department of Arms and Armor at the Metropolitan Museum of Art in New York, where he amassed an enormous collection in this field.

Thanks to gifts and subscriptions from Dr. Dean's family, friends and colleagues, the committee was successful in securing funds for the reproduction of the plates while the trustees of the American Museum of Natural History undertook to publish the Dean Memorial Volume on Archaic Fishes as fast as the separate parts could be completed by several specialists.

Under the able editorship of Dr. Eugene Willis Gudger four parts of the volume have already appeared while several others are in press or otherwise in progress.

In Article I, A Biographical Sketch of Bashford Dean by the present writer, are set forth some of the home influences and educational experiences that led to Dean's delightful personality and unique achievements.

Article II, "The Segmentation of the Egg of the Myxinoid, *Bdellostoma stouti*, Based on the Drawings of the Late Bashford Dean," by E. W. Gudger and Bertram

G. Smith, describes the earlier stages of cleavage of the egg of *Bdellostoma stouti*, the latter stages having been described in Dean's own memoir of 1899.

In Article III, "The Genital System of the Myxinoidea: A Study based on Notes and Drawings of these Organs in *Bdellostoma* made by Bashford Dean," Dr. J. LeRoy Conel confirms Dean's finding that *Bdellostoma stouti* is certainly not hermaphroditic in any respect, even although many individuals of *Myxine* have either a functional testis and a non-functional ovary or the reverse, as shown by A. and K. E. Schreiner in 1905 and 1908, in continuation of Dean's earlier inquiries on this subject.

Article IV by Anatol Heintz is entitled "The Structure of *Dinichthys* A Contribution to our Knowledge of the Arthrodira."

Dinichthys was an extinct type of giant fish-like vertebrates the fossil remains of which are found in the Cleveland Shale, of Upper Devonian Age. The big skull and jaw plates early engaged the attention of collectors, the first reconstruction of the skull and dorsal carapace having been published in 1875 by the geologist Newberry, Dean's professor at Columbia University. The scientific game of fitting these scattered pieces together was played by Newberry, Claypole, Dean and many of their successors, but it has remained for the young curator of the Paleontological Museum at Oslo, Norway, after a year of intensive investigation of the collections left by Newberry and Dean, to evolve a new and definitive restoration of the skeleton of *Dinichthys*. By acquiring an irrefutable knowledge of the contacts of every bone on this three dimensional picture puzzle Dr. Heintz has not only succeeded in detecting radical errors in all earlier restorations but has also paved the way for his far-reaching conclusions concerning the relationships of the extinct group of Arthrodira to which *Dinichthys* belongs.

More in detail some of the outstanding features of this contribution are as follows:

(1) The pair of joints between the back of the head and neck, the shield to which the group name Arthrodira refers, acted like a pair of hinges on a gate except that they were in a horizontal plane. Owing to the vertical crushing of the convex plates that carried the cylinder and the socket respectively, the opposite hinge joints are usually pitched downward in the actual fossils, but Heintz shows that they could only have functioned if they were horizontal in life and that when the convexity of the intervening portion of the plates is properly restored the joints permit the raising of the head on the neck plates. All previously published restorations had been at fault in this respect. When this critically important correction is made the whole shape of the head becomes rounder and more fish-like.

(2) Heintz confirms and extends the general interpretation of the mechanism and movements of the head and mandible which had been independently proposed by Jaekel and by L. A. Adams in 1919. He concludes with Adams that there were four sets of muscles involved in this movement. This arrangement is entirely unique among fish-like vertebrates outside of the Arthrodira and its presence supports Dr. Heintz's conclusion that the Arthrodira were widely removed from both the Elasmobranchs and the higher fishes. At the same time he confirms the general conclusion of Dean and of Hussakof that the Arthrodira were related to the Antiarchi and together constituted an independent extinct class of vertebrates, the Placodermata of McCoy. We may welcome both the definitive publication on *Dinichthys* and the establishment of a landmark in the early evolution of the vertebrates.—WILLIAM K. GREGORY, *American Museum of Natural History, New York City.*

EDITORIAL NOTES AND NEWS

Piscium Catalogus

THE German scientific publisher W. JUNK (Berlin W. 15, Sächsische Str. 68, has just announced the proposed publication, beginning with January 1933, of *PISCIMUM CATALOGUS*, a synonymic and distributional check-list of the fishes of the world, to be accompanied by a complete bibliography and index. The literature on the economically important species would be included. The competence of the firm to handle this large project is attested by the fact that it is now publishing three similar catalogs, excellently covering Coleoptera, Lepidoptera and fossils. More than 240 parts of these three catalogs have appeared.

About 16 volumes of 400 pages each are anticipated, to cover the estimated 500 families and subfamilies, 5000 genera and 40000 species of fishes.

The crying need for a work of this sort is evident to every ichthyologist. The last world treatise on fishes was Günther's *Catalogue of the Fishes of the British Museum*, published 62 to 73 years ago. Most of the advance in systematic ichthyology has been made in the last half-century. Günther treated approximately only one-fifth the number of genera and species now recognized. Not only is Günther's *Catalog* almost hopelessly out-of-date, but it is almost impossible to obtain. The eight volumes now sell for more than \$200.00.

The publication of this monumental work is dependent on the number of advance subscriptions received. The price as announced is Mk. 1.60 per signature, and the annual cost is estimated as about 48 Mk.—. The work should prove virtually indispensable in all major ichthyological research, should be in the hands of every working ichthyologist and should be on the shelves of every zoological museum and of every library devoted wholly or in part to natural science.

If the *Piscium Catalogus* can be undertaken, Junk proposes to publish also an *AMPHIBIORUM ET REPTILIORUM CATALOGUS*.

New York Zoological Society

THE New York Zoological Society has remodelled a one-story brick building in its Zoological Park, to serve as the laboratory for the Department of Tropical Research, and as a museum to hold the varied collections made by DR. WILLIAM BEEBE and his associates.

At the New York Aquarium, on November 2, DR. CHARLES HASKINS TOWNSEND was honored at a reception by the staff and employees of the Aquarium. The occasion was the completion of thirty years in the directorship of the institution.

Herpetological Items

DR. E. R. DUNN with the able assistance of Mrs. Dunn and of his student, Jean Piatt, is engaged in a revision of the herpetological collections of the Philadelphia Academy of Natural Sciences.

The Swiss herpetologist, DR. RUDOLF STÖHLER, and Mrs. Stöhler are living this year in the United States. He has been spending some time in the museums of the country continuing his observations on the sex organs of the Bufonidae, and is now in Oakland, California (3537 Galindo St.).

Word just received from KARL P. SCHMIDT at Vienna states that he is completing his round of European museums, and will sail for America on January 6.

The herpetological material collected by Major Chapman Grant during the past few years on Porto Rico and adjacent islands, comprising over seven thousand specimens including his types, and beautifully representing the herpetological fauna of these islands, has been acquired by the Museum of Zoology of the University of Michigan and the Museum of Comparative Zoology of Harvard University. Major Grant is now stationed at Indianapolis, Indiana (412 East Fall Creek Boulevard).

**1933
Meeting**

THE American Society of Ichthyologists and Herpetologists will hold its annual meeting for 1933 at the Museum of Comparative Zoology, Harvard University, Cambridge, Massachusetts, in May. As for the several years past, the meeting will be held in conjunction with the annual gathering of the American Society of Mammologists. A further announcement will either be included in the next number of COPEIA or sent through the mail.

An interesting meeting is in prospect. The Secretary will be glad to receive word from those who expect to attend, and titles of the papers they may contribute.

**National
Museum**

DR. GEORGE S. MYERS, of Stanford University, has just been appointed Associate Curator of Fishes in the U. S. National Museum. The Editors of COPEIA wish to compliment the Smithsonian authorities for filling this important position at the present time, and to congratulate Dr. Myers on the appointment, for which he is admirably fitted.

**The National
Aquarium
Society**

THE newly organized National Aquarium Society held its first regular meeting in the Aquarium of the Commerce Building on November 18, 1932, with an attendance of about 100 persons including members and visitors. The officers of the society were greatly pleased over the enthusiasm and interest taken. Twenty persons submitted their names to become members. The object of the society shall be the popular and scientific study of the aquarium, its flora and fauna.

The officers of the society consist of J. J. FITZPATRICK, president; WILLIAM BROWN, vice president; and a board of councilors consisting of DR. LEWIS RADCLIFFE, DR. PAUL BARTSCH, DR. WILLIAM MANN, and FRED G. ORSINGER.

It is the intention of the society to hold regular monthly meetings throughout the year with the exception of July and August.—From *Fisheries Service Bulletin*.

**The Plural
of Fish**

MANY questions are asked as to the proper plural of the word "fish." Answers differ. The editorial policy of COPEIA is to use "fish" as a collective plural when more than one *individual* is referred to, but to use "fishes" when more than one *species* is indicated. This is in line with common practice in forming the plural of other animal names, as, for example, "deer."

**Recent
Death**

DR. WILLIAM PATTEN, Professor Emeritus of zoology at Dartmouth College, widely known for his theory that the vertebrates evolved from specialized arachnids, and for his researches on fossil arachnids and fossil fishes, died on October 27, at the age of seventy-one.

INDEX FOR 1932

New names and their principal page references are printed in bold face type

- abacura, Farancia 37
 Aboma chiquita 168
 Abudeiduf declivifrons 167
 marginatus 167
 acanthias, Squalus 31, 181
 Acanthodactylus 73
 Acanthophaelus reticulatus 33, 34
 Acentrophorus 54
 Acheilognathus mesembrinum 176
 Achirus fasciatus 32
 Acrotus willoughbyi 65
 Actinemys marmorata 9
 aculeatus, Polypedates 179
 acutus, Fodiator 164
 adobe, Apocope 135
 Aedes 84
 aeneus, Aneides 101, 102
 Myoxocephalus 32
 aequidens, Cuius 69
 aestivus, Ophedryx 7, 8, 37
 agassizii, Gopherus 128
 Agkistrodon mokasen 38
 Aglyptocephalus madagascariensis 178
 albolineatus, Brachydania 23
 Albula vulpes 164
 aleuticus, Cottus 19
 Alexander, Gordon 78
 aliata, Corynopoma 33
 Stewardia 33
 alleganiensis, Cryptobranchus 174
 Allen, Morrow J. 104, 105, 179
 alleni, Liodytes 37
 Allopoecilia caucana 138
 almadensis, Leimadophis 89, 92
 Alsophis 89, 90, 91
 angulifer 90
 anomalus 90
 antillensis 92
 ater 90
 melanichnus 90
 alvarius, Bufo 175, 176
 Amastridium 91
 ambinigra, Apostolepis 103
 amblystomopsis, Cottus 17, 19, 20
 Ambystoma maculatum 2, 84
 tigrinum 98
 americana, Morone 22
 americanus, Ammodytes 31
 Bufo 134
 Bufo lentiginosus 134
 Pseudopleuronectes 32
 Ammodytes americanus 31
 Amphisbaena darwini 178
 occidentalis 178
 townsendi 178
 andersoni, Hyla 42
 Aneides 101, 102
 aeneus 101, 102
 Angiostomum 25
 Anguilla 26
 anguineus, Chlamydoselachus 182
 angulifer, Alsophis 90
 Anisotremus interruptus 166
 annectens, Pituophis catenifer 125
 Anniella 4, 77
 nigra 5, 6
 pulchra 4, 5, 6
 annulatus, Sargus 24
 Spheroides 167
 Anolis 11
 baccatus 12
 utowanae 12
 Anomalepis 8
 mexicanus 7
 anomalus, Alsophis 90
 Anopiarachus 143, 144, 146
 purpurescens purpurescens 143
 ansatus, Trachidermus 20
 Antennarius 60
 Antigonia 58
 antillensis, Alsophis 92
 Apeltes 33
 quadracus 33
 Apocope 135
 adobe 135
 carringtonii 135
 couesii 135
 oscula 135
 oscula 135
 jarrowi 135
 Apostolepis ambinigra 103
 dorbignyi 103
 tenuis 103, 104
 appendiculatus, Polypedates 179
 approximans, Holbrookia maculata 103, 176
 arenicolor, Hyla 176
 argentatus, Clinus 146
 argentea, Sphyræna 165
 argenteum, Hyperprosopon 167
 Argentina 105
 sialis 105
 Arizona elegans occidentalis 125
 Armstrong, J. C. 33
 Arrhyton 89, 90, 91, 92
 taeniatum 90
 vittatum 90
 Ascaphus 100
 truei 38, 100
 Asellus 84, 85
 Astroblepus 137
 festae 138
 fissidens 138
 grixalvii 137

heterodon	138
longifilis	137
nicefori	137
peruanus	138
simonsi	138
Astroscoptes	58
Astyanax fasciatus	137
integer	137
ater, Alsophis	90
Dromicus	89, 90
Atherinops affinis magdalenae	165
Atherinopsis sonorae	165
atlanticus, Tarpon	28
atrata, Nimia	6
atropurpureus, Epigeeichthys	143
atrovirens, Sebastodes	167
aubryi, Leptopelis	179
auratus, Chrysophrys	24
Mullus	180
Aurelia sp.	180
aurifasciatus, Philautus	179
australis, Clinocottus analis	168
azalea, Runula	168

Babcock, Harold L.	43
baileyi, Crotaphytus collaris	120
Balsadichthys	68
Barbodes disa	140
sirang	140, 141
Barbour, Thomas	11, 97
baroni, Mantella	179
Bathygobius soporator	168
Bdellostoma stouti	182, 183
bellii, Chrysemys	9, 10
Chrysemys picta	9, 10
Emys	9
Belostoma	84
Berg, Leo S.	17, 156, 159
bilinearis, Merluccius	32
bimaculatus, Polypedates	179
biseriatus, Sceloporus occidentalis	122
Bishop, Sherman C.	1
bislineata, Eurycea bislineata	84, 86
blainvillii, Phrynosoma blainvillii	123
Blanchard, Frank N.	98
Frieda Cobb	98
blandingii, Emys	43, 173, 174
Blennius gattorugina	146
gunnellus	143, 144, 146
montagui	146
ocellarius	145
pholis	144, 145
sphinx	146
bocagii, Leptopelis	179
borealis, Sphyræna	31
Box	24
boylly, Lampropeltis getulus	98, 126
braccatus, Anolis	12
Brachydanio albolineatus	23
rerio	23
brachynemus, Megalixalus	179
Breder, Jr., C. M.	32, 147, 180
brevifolia, Yucca	123
Brevigobio kawabatae	156
brevirostris, Leptopelis	179

Brochymena sulcata	14
Bufo	78
alvarius	175, 176
americanus	134
copei	134
californicus	119
canagicus halophilus	119
cognatus	119
cognatus	175
copei	134
lentiginosus americanus	134
melanostictus	78, 80
valliceps	158
vulgaris	79
Burnett, W. L.	37
Burt, Charles E.	104, 158, 181
butleri, Thamnophis	113, 114, 115, 116, 117
Thamnophis radix	96
Butorides virescens	37

Caecilia intermedia	6
caerulea, Sardinops	65, 164
Cahn, Alvin R.	86
californica, Squatina	65
californicus, Bufo	119
Myliobatis	163
californiensis, Eucinostomus	166
Xenistius	166
callarias, Gadus	22
Callechelys myersi	139
melanotaenia	139
callilaemus, Dromicus	89, 90
Callionymus	58
Callisaurus	72, 73, 76, 77
crinitus	73
ventralis	72
ventralis	121, 176
Cambarus	84
canis, Mustelus	31
Carcharias lamia	50
littoralis	31, 48
carolina, Terrapene carolina	86
carolinus, Prionotus	32
carringtonii, Apocope	135
Castanopsis sempervirens	36
catesbyi, Uromacer	91
caucana, Allopoecilia	138
Caularchus maeandricus	143
Caulolatilus princeps	168
Cebidichthys violaceus	143
Centropristes	31, 32
striatus	32
centrura, Dasyatis	31
cepedianum, Dorosoma	34
Cerastes	73, 74, 75
cornuta	73
vipera	73
cerastes, Crotalus	4, 74, 127
Cestracion philippi	182
Cetengraulis mysticetus	64
Chaenomugil proboscideus	165
Chaetodipterus	58
Chaetodon ocellatus	180
Chalcides sepoides	75, 76, 77, 78

- chalybaeus, *Geophis* 8
 chamissonis, *Dromicus* 90
 Leimadophis 92
 Characodon 68
 furcoides 68
 Charax 24
 Chatoessus punctatus 30
 Cheirolepis 53, 55
 Chelinidea vittiger 14
 Chelonia mydas 43
 Chelydra serpentina 37
 chilensis, *Sarda* 165
 Chilomeniscus cinctus 177
 chiquita, *Aboma* 168
 Chiromantis petersii 179
 rufescens 179
 xerampelina 179
 Chlamydoselachus anguineus 182
 chlorostictus, *Sebastodes* 168
 Chorinemus 156
 Chrysemys bellii 9, 10
 marginata 9
 picta 37
 bellii 9, 10
 marginata 43
 chrysochloris, *Pomolobus* 32
 Chrysophrys auratus 24
 chrysops, *Stenotomus* 22, 23, 24, 25, 32
 chuss, *Phycis* 32
 cinctus, *Chilomeniscus* 177
 cinereum, *Xystaema* 166
 cinereus, *Plethodon* 85
 Citharichthys fragilis 168
 clarkii, *Notolepidomyzon* 136
 Sceloporus 176
 clathratus, *Paralabrax* 166
 Clemmys decussata 132
 guttata 37
 marmorata 9
 muhlenbergii 174
 Clinocottus analis australis 168
 Clinus argentatus 146
 Clupea harengus 31
 Cnemidophorus 73, 121
 sexlineatus perplexus 158, 176
 sackii 158
 tessellatus 122
 tessellatus 123
 Coates, C. W. 147
 cognatus, *Bufo* 119
 Bufo cognatus 175
 Coleonyx variegatus 120
 colias, *Pneumatophorus* 31
 Cololabis saira 66, 67
 Coluber constrictor mormon 177
 completa, *Ctenosaura* 7
 compressicauda, *Natrix* 89
 concolor, *Hyperolius* 179
 Conder, Evert 86
 conger, *Leptocephalus* 31
 Coniophanes 7, 8, 91
 consobrinus, *Sceloporus* 99
 Sceloporus undulatus 158
 constellatus, *Sebastodes* 168
 Contia 91
 copei, *Bufo* 134
 Bufo americanus 134
 corallinus, *Micrurus* 6, 8
 Coregonus coulterii 62
 oregonius 63
 Corixa 84
 cornuta, *Cerastes* 73
 Corydoras melanotaenia 137
 Corynopoma 33
 aliata 33
 riisei 33
 Cottus 19
 aleuticus 19
 amblystomopsis 17, 19, 20
 czerskii 17, 19, 20
 emeljanovi 19
 hangiongensis 19
 kaganowskii 17, 18, 19
 kasika 19
 knerii 19
 paltschevskii 19
 poecilopus 17
 scorpius 24
 couchii, *Scaphiopus* 175
 couesii, *Apocope* 135
 coulterii, *Coregonus* 62
 Prosopium 62, 63, 64
 Creaser, Charles W. 157
 crinitus, *Callisaurus* 73
 Crotalopsis 105
 punctifer 105
 Crotalus 74
 cerastes 4, 73, 74, 127
 confluentus lutosus 127
 mitchellii 127
 oreganus 36, 127, 169, 170, 171
 stephensi 127
 durissus 171
 horridus 171
 molossus 38
 ruber 38
 scutulatus 119, 127
 Crotaphytus collaris baileyi 120
 wislizenii 121, 122
 crumenophthalmus, *Trachurops* 165
 Cryptobranchus 3
 alleganiensis 174
 Ctenosaura 8, 97
 completa 7
 cuitzeoensis, *Zoogoneticus* 69
 Culius aequidens 69
 Curimatus metae 137
 cursor, *Dromicus* 92
 cuvieri, *Scolecossaurus* 7
 cyanogenys, *Sceloporus torquatus* 158
 cyclopion, *Natrix* 37
 Cyclops 84
 Cyclopterus lumpus 32
 Cynoscion parvipinnis 167
 regalis 32
 czerskii, *Cottus* 17, 19, 20
 danielsi, *Gyrinophilus* 81, 82, 83
 darwini, *Amphisbaena* 178
 Dasyatis centrura 31

- .134
.134
.6, 8
.62
.63
.84
.73
.137
.33
.33
.33
.19
.19
.20
.20
.19
.19
.19
.19
.19
.19
.24
.175
.135
.62
.64
.157
.73
.105
.105
.74
.127
.127
.127
.171
.127
.171
.38
.38
.127
.120
.165
.3
.174
.97
.7
.69
.69
.137
.92
.7
.158
.37
.84
.32
.167
.32
.20
.83
.78
.31
- dipterura163
Davis, D. Dwight113
declivifrons, Abudehduf167
decurrens, Pleuronichthys67
decurtatus, Phyllorhynchus119, 125
decussata, Clemmys132
Emys132
Pseudemys129, 131, 132, 133
dekayi, Storeria7
De Lacy, Allan C.143
delphinus, Notolepidomyzon136
dentatus, Paralichthys32
deserticola, Pituophis catenifer125
Desmognathus84, 85, 95
fuscus fuscus86, 101
ochrophaeus101
phoca101
De Sola, C. Ralph45, 129
Diadophis8
punctatus edwardsii101
punctatus7, 8
strictogenys7, 8
diaphanes, Raja31
diaphanum, Ostracion181
Diaptomus84
diazi, Zoogoneticus69
Dicamptodon ensatus38
diego, Pneumatophorus165
Dinichthys183
Dipodomys73
Dipsosaurus73, 122
dorsalis dorsalis4, 120
dipterura, Dasyatis163
Dipus73
disa, Barbodes140
dorbignyi, Apostolepis103
Dormitator latifrons68
maculatus69, 168
Dorosoma cepedianum34
dorsalis, Dipsosaurus dorsalis4, 120
Megalixalus179
dovii, Gymnothorax164
draytonii, Rana aurora120
Dromicus89, 90, 91
ater89, 90
callilaemus89, 90
chamissonis90
cursor92
ferox90
parvifrons protenus91
stahli90
ductor, Naucrates32
dugessii, Zoogoneticus69
Dunn, E. R.89, 97
dunni, Geophis7, 8
durissus, Crotalus171
duryi, Gyrinophilus81, 82, 83
Pseudotriton81
Dymond, John R.31, 102, 182
Dytiscus84
ebenau, Mantella179
Echeneis51
naucrates45, 48, 49, 50, 52
edwardsii, Diadophis punctatus101
eglanteria, Raja31
elegans, Micrurus7
Uta stansburiana176
Virginia valeriae7
Eleotris picta68
Eleutherodactylus97
nubicola97
ranoides emleni97
nubicola97
elongatus, Sceloporus14
Sebastodes167
emeljanovi, Cottus19
emleni, Eleutherodactylus ranoides9
Emys bellii9
blandingii173, 174
decussata132
oregonensis9
rugosa132
Engelhardt, Geo. P.38
Engraulis mordax mordax164
ensatus, Dicamptodon38
Epigeichthys atropurpureus143
eques, Polypedates179
Eretmochelys imbricata48
erinacea, Raja31
Etumeus micropus164
Eubbranchipus84
Eucinostomus californiensis166
Eumeces fasciatus37, 104
multivirgatus104
obsoletus99
Eurycea bislineata bislineata84, 86
longicauda94
euryops, Thorophos61
euryxanthus, Micruroides6
Evermannella162
Evermannia162
Evermannichthys162
Evermanolus162
Ewan, Joseph36
exilis, Strongylura164
eximia, Hyla99
Farancia abacura37
fario, Salmo21
fasciatus, Achirus32
Astyanax137
Eumeces37, 104
Trachidermus17, 20
ferox, Dromicus90
Hypsirhynchus90
festae, Astroblepus138
fissidens, Astroblepus138
Fistularia58
flavescens, Perca21
floridana, Pseudemys92, 93
Fodiator acutus164
forasini, Megalixalus179
formosus, Sclerophages30
fragilis, Citharichthys168
frenatus, Masticophis flagellum124, 128, 177
Fry, F. E. J.102
fucorum, Xerepes143
fulvius, Micrurus7

- Micrurus fulvius* 6
Fundulus heteroclitus macrolepidotus 31
 majalis 23, 31
furcoides, *Characodon* 68
fuscus, *Desmognathus fuscus* 86, 101
 Syngnathus 180

Gadus callarias 22, 32
Gaige, *Helen T.* 107, 134
gairdnerii, *Salmo* 32
galcoides, *Otophidium* 168
Gammarus 84
Gastrostomus 58
gattorugina, *Blennius* 146
Genyonemus lineatus 105, 167
geographica, *Graptemys* 174
Geophis 6, 7, 8
 chalybaeus 8
 dunni 7, 8
 semidoliatus 7, 8
Gerres lineatus 166
getulus, *Lamprozelis* 98
giganteus, *Saxidomus* 144
gilberti, *Hypsoblennius* 168
Girella nigricans 166
gladius, *Xiphias* 32
glandulosus, *Philautes* 179
glaucia, *Prionace* 31
glaucum, *Stizostedion* 182
glutinosus, *Plethodon* 85, 102
Gobiesox 60
Gobiosoma histrio 168
 molestum 28
Goff, *C. C.* 92
 Dorothy S. 92
Goodea whitei 68
goodei, *Sebastodes* 167
Gopherus agassizii 128
goudoti, *Polypedates* 179
graciosus, *Sceloporus graciosus* 13
grandiflora, *Sesbania* 97
Graptemys geographica 174
Greenhall, *Arthur* 129
Gregory, *William K.* 53, 103
grixaivii, *Astroblepus* 137
Gunderson, *Fred J.* 65
gunnellus, *Blennius* 143, 144, 146
 Pholis 144, 146
guttata, *Chrysemys* 37
guttatus, *Lutianus* 166
guttulatus, *Mantidactylus* 179
Gymnothorax dovii 164
Gyrinophilus 81, 82, 95
 danielsi 81, 82, 83
 duryi 81, 82, 83
 porphyriticus 81, 82, 83, 85, 101

haitej, *Mesocottus* 17, 19, 20
Halipilus 84, 85
halophilus, *Bufo canagicus* 119
Hamilton, Jr., *William J.* 83
hammondii, *Scaphiopus* 36, 104, 119, 175
hangiongensis, *Cottus* 19
Hanna, *G. D.* 161
harengus, *Clupea* 31

Hart, *John Lawson* 65
Hartweg, *Norman* 103
Hassler, *William G.* 94
Hatt, *Robert T.* 37
Helminthophis 7, 8
Heloderma suspectum 177
Hemianthias peruanus 106
Hemigrammocypris 156
 rasborella 156
Hemitripterus 54
hernandesi, *Phrynosoma douglasii* 177
Herpele 8
 ochrocephala 7
Herre, *Albert W. C. T.* 139
Heterodon 72
heterodon, *Astroblepus* 138
Hexagenia 84
hexalepis, *Salvadora grahamiae* 124
Hill, *Howard R.* 181
Histrio 60
 histrio, *Gobiosoma* 168
Hobbs, *Kenneth L.* 104
Holbrookia 78, 103
 maculata approximans 103, 176
 maculata 103
 texana 176
holbrookii, *Lamprozelis getulus* 98
 Scaphiopus 35
holbrookii, *Scaphiopus* 104
Hopkins, *C. E. P.* 157
horridus, *Crotalus* 171
horstocki, *Hyperolius horstocki* 179
Hubbs, *Carl L.* 26, 68, 105, 107, 159
humilis, *Leptotyphlops humilis* 4
Hyalala 84
Hybopsis 74
Hyborrhynchus 84
Hyla 99
 andersoni 42
 arenicolor 176
 eximia 99
 regilla 104, 120
Hylambates hylodes 179
 maculatus 179
 natalensis 179
hylodes, *Hylambates* 179
Hynobius ikishimae 158
 nebulosus 158
 tagoi 158
 tsuensis 158
Hyperolius concolor 179
 horstocki horstocki 179
 kivuensis 179
 marginatus 179
 mariae 179
 marmoratus 179
 montana 179
 ocellatus 179
 platyrhinus 179
 picturatus 179
 puncticulatus 179
 rhodoscelsis 179
 rossii 179
Hyperprosopon argenteum 167
Hyporhamphus roberti 164, 180

.. 65	Hypsigena	91	bocagii	179
.. 103	ochrorhynchus	103, 126	brevirostris	179
.. 94	Hypsirhynchus	89, 90	johnstoni	179
.. 37	ferox	90	notatus	179
.. 7, 8	Hypsoblennius gilberti	168	ocellatus	179
.. 177			palmaris	179
.. 106	Ialtris	89, 90, 91, 92	rufus	179
.. 156	ikishimae, Hynobius	158	uluguruensis	179
.. 156	imbricata, Eretmochelys	48	Leptotyphlops humilis humilis	4
.. 54	imperialis, Luvarus	181	lepturus, Trichiurus	180
.. 177	infernalis, Thamnophis sirtalis	127	leucomystax, Polypedates	179
.. 8	integer, Astyanax	137	leucorhinus, Philautus	179
.. 7	intermedia, Caecilia	6	Lichanura roseofusca roseofusca	124
.. 139	interruptus, Anisotremus	166	linearis, Uta ornata	176
.. 72	intertinctus, Mystriophis	105	lineatus, Genyonemus	167
.. 138	Irillion	63	Gerres	166
.. 84			Katsuwonus	165
.. 124	jarrovi, Sceloporus	177	Limnaea	84
.. 181	Johnson, H. Herbert	21	Limnophilus	84
.. 60	johnstoni, Leptopelis	179	Liodytes alleni	37
.. 168			Liopeltis vernalis	102
.. 104	kaganowskii, Cottus	17, 18, 19	Liophis	91
.. 103	Kassina senegalensis	179	littoralis, Carcharias	31, 48
.. 176	Katsuwonus lineatus	165	longicauda, Eurycea	94
.. 103	pelamis	165	longifilis, Astroblepus	137
.. 176	kawabatae, Brevigobio	156	Lophius	60
.. 98	kazika, Cottus	19	Lophopsetta maculata	32
.. 35	Kellogg, Remington	36	Loxolophodon	40
.. 104	King, F. Willis	99, 175	Lumpus, Cyclopterus	32
.. 157	Kinosternon subrubrum subrubrum	43	Lutianus	163
.. 171	kisutch, Oncorhynchus	21	guttatus	166
.. 179	kiyuensis, Hyperolius	179	lutosus, Crotalus confluentus	127
.. 159	Klauber, L. M.	4, 100, 118	Luvarus imperialis	181
.. 4	knerii, Cottus	19		
.. 84	Konosirus	30	macrolepidotus, Fundulus heteroclitus	31
.. 74			macropterus, Neothunnus	165
.. 84	Labrisomus xanti	168	macrotris, Polypedates	179
.. 99	lacertina, Siren	37	maculata, Holbrookia maculata	103
.. 42	laetus, Pholis	143	Lophopsetta	32
.. 176	lamia, Carcharias	50	Ophis	26, 27
.. 99	La Monte, Francesca	34	maculatum, Ambystoma	2, 84
.. 120	Lampropeltis getulus	98	maculatus, Dormitator	69, 168
.. 179	boylii	98, 126	Hylambates	179
.. 179	holbrookii	98	Philypnus	68
.. 179	triangulum triangulum	98	Polypedates	179
.. 179	laterale, Leiopisma	7	Spheroides	32
.. 158	lateralis, Masticophis	124	Zoogoneticus	69
.. 158	Philypnus	69	maculosa, Ophis	27
.. 158	latifrons, Dormitator	68	maculosus, Necturus	1, 83
.. 158	Leavitt, Benjamin	179	Oligocottus	143
.. 179	Lebistes	23, 147, 151, 153	madagascariensis, Aglyptocephalus	178
.. 179	reticulatus	33, 147	Megalixalus	179
.. 179	lecontei, Rhinocheilus	126	maeandricus, Caularchus	143
.. 179	Leimadophis	89, 92	magdalenae, Atherinops affinis	165
.. 179	almadensis	92	magdalenae, Micrognathus	141
.. 179	chamissonis	92	magister, Sceloporus	103, 122
.. 179	Leiopisma	8	majalis, Fundulus	23, 31
.. 179	laterale	7	Mantella baroni	179
.. 179	Lepidomeda vittata	135	ebenaui	179
.. 179	Lepidosteus	54	Mantidactylus guttulatus	179
.. 179	Leporinus y-ophorus	137	ulcerosus	179
.. 179	Leptocalamus	91	marginata, Chrysemys	9
.. 179	Leptocephalus conger	31	Chrysemys picta	43
.. 167	Leptodeira	91	marginatus, Abudedefduf	167
.. 80	Leptopelis aubryi	179	Hyperolius	179

- mariae, *Hyperolius* 179
 marina, *Strongylura* 180
 marinus, *Petromyzon* 32, 157
 marmorata, *Actinemys* 9
 Clemmys 9
 Ulca 67
 marmoratus, *Hyperolius* 179
 Synbranchus 7
 Masticophis flagellum frenatus 124, 128, 177
 lateralis 124
 matafae, *Micrognathus* 142
 m'callii, *Phrynosoma* 78, 100
 mecullochi, *Pseudochromus* 30
 Megalixalus brachynemis 179
 dorsalis 179
 forasini 179
 madagascariensis 179
 megalops, *Thamnophis* 35
 melanichnus, *Alsophis* 90
 melanostictus, *Bufo* 78, 80
 melanotaenia, *Callechelys* 139
 Corydoras 137
 Menidia menidia notata 31
 Menticirrhus undulatus 167
 Merluccius bilinearis 32
 mesembrina, *Metzia* 156
 mesembrinum, *Acheilognathus* 156
 Mesocottus haitej 17, 19, 20
 metae, *Curimatus* 137
 Metzia 156
 mesembrina 156
 mexicanus, *Anomalepis* 7
 Micrognathus magdamoi 141
 matafae 142
 negrosensis 142
 micropus, *Etrumeus* 164
 Micruroides euryxanthus 6
 Micrurus 6
 corallinus 6, 8
 riisei 7
 elegans 7
 fulvius 7
 fulvius 6
 mipartitus 7
 nigrocinctus nigrocinctus 7, 8
 Mimometopon 91
 mipartitus, *Micrurus* 7
 mitchellii, *Crotalus confluentus* 127
 mocquardi, *Polypedates* 179
 mohavensis, *Yucca* 123
 mokasen, *Agkistrodon* 38
 Mola mola 180
 molestum, *Gobiosoma* 28
 molossus, *Crotalus* 38
 montagi, *Blennius* 146
 montana, *Hyperolius* 179
 montanus, *Pseudotriton* 81
 mordax, *Engraulis mordax* 164
 Osmerus 157
 mormon, *Coluber constrictor* 177
 Morone americana 22
 Mosauer, Walter 72
 Mugil cephalus 165
 curema 165
 muhlenbergii, *Clemmys* 174
 Mullus auratus 180
 multivirgatus, *Eumeces* 104
 mundus, *Oligoplites* 165
 Muraena ophis 27
 Mustelus canis 31
 Mycteroperca 163
 xenarcha 165, 166
 mydas, *Chelonia* 43
 Myers, George S. 30, 35, 39, 41, 61, 62, 137
 myersi, *Callechelys* 139
 Myliobatis californicus 163
 Myoxocephalus 54
 aeneus 32
 octodecemspinosus 32
 scorpius 24
 Myrophis vafer 164
 mysticetus, *Cetengraulis* 164
 Mystriophis 105
 intertinctus 105
 Myxine 183
 natalensis, *Hylambates* 179
 Natrx 89, 92
 clarkii 179
 compressicauda 89
 cyclopion 37
 Naucrates ductor 32
 naucrates, *Echeneis* 45, 48, 49, 50, 52
 Nealosa 30
 nebulifer, *Paralabrax* 166
 nebulosus, *Hynobius* 158
 Percis 58
 Necturus 1, 3, 83, 84
 maculosus 1, 83
 negrosensis, *Micrognathus* 142
 Neophos 61
 nexilis 61
 Neothunnus macropterus 165
 Nesiotas 30
 purpurascens 30
 Netting, M. Graham 101, 173
 nexilis, *Neophos* 61
 niceforii, *Astroblepus* 137
 nigra, *Anniella* 5, 6
 nigricans, *Girella* 166
 nigriceps, *Tantilla* 177
 nigrocinctus, *Micrurus nigrocinctus* 7, 8
 Ninia 8
 atrata 6
 sebae 7
 notata, *Menidia menidia* 31
 Uma 73, 75, 122
 notatus, *Leptopelis* 179
 Notolepidomyzon 136
 clarkii 136
 delphinus 136
 santa-anae 136
 utahensis 135, 136
 Notonecta 84
 nubicola, *Eleutherodactylus* 97
 Eleutherodactylus ranoides 97
 obesus, *Sauromalus* 121

- .174
.180
.104
.165
.27
.31
.163
.166
.43
.137
.139
.163
.54
.32
.32
.24
.164
.164
.105
.105
.183
.179
.92
.179
.89
.37
.32
.0, 52
.30
.166
.158
.58
.84
.83
.142
.61
.61
.163
.30
.30
.173
.61
.137
.5, 6
.166
.177
.7, 8
.8
.6
.7
.31
.122
.136
.136
.136
.136
.84
.97
.97
.121
- obsoletus, Eumeces 99
occidentalis, Amphisbaena 178
 Arizona elegans 125
 occipitalis, Sonora 4, 77, 126
 ocellarius, Blennius 145
 ocellatus, Chaetodon 180
 Hyperolius 179
 Leptopelis 179
 ochrocephala, Herpele 7
 ochrophaeus, Desmognathus fuscus 101
 ochrorhynchus, Hypsiglena 103, 126
 octodecemspinosus, Myoxocephalus 32
 officinalis, Scincus 76
 Ogocephalus 60
 Oligocottus maculosus 143
 Oligoplites mundus 165
 Oncorhynchus kisutch 21
 onitis, Tautoga 32
 Ophisaurus 8
 aestivus 7, 8, 37
 Ophichthus 27
 Ophis 26, 27
 maculata 26, 27
 maculosa 27
 ophis, Muraena 27
 Ophisaurus 8
 ventralis 7
 Opsanus tau 32, 60
 oreganus, Crotalus confluentus 36, 127, 169, 170, 171
 oregonensis, Emys 9
 oregonium, Prosopium 63
 oregonius, Coregonus 63
 ornata, Terrapene 177
 ornatus, Pholis 143
 oscula, Apocope 135
 Apocope oscula 135
 Osmerus mordax 157
 Osphronemus 180
 saigonensis 180
 siamensis 180
 ostoni, Polypedates 179
 Ostracion diaphanum 181
 Otophidiu galeoides 168
 Oyama, Junji 158
 pachycephalus, Zoogoneticus 69
 Pagellus 24
 palmatus, Leptopelis 179
 palustris, Pseudemys 129
 Paralabrax clathratus 166
 nebulifer 166
 Paralichthys dentatus 32
 pardalis, Polypedates 179
 Parker, H. W. 178
 parvipinnis, Cynoscion 167
 pelamis, Katsuwonus 165
 Perca flavescens 21
 Percis 58
 nebulosus 58
 perplexus, Cnemidophorus sexlineatus 158, 176
 peruanus, Astroblepus 138
 Hemianthias 106
 petersii, Chiromantis 179
- Petromyzon marinus 32, 156
Philautus aurifasciatus 179
 glandulosus 179
 leucorhinus 179
 vittatus 179
philippi, Cestracion 182
Philypnus lateralis 69
 maculatus 68
phoca, Desmognathus 101
Pholas 144
Pholis gunnellus 144, 146
 laetus 143
 ornatus 143
pholis, Blennius 144, 145
Phrynocephalus 73
Phrynosoma 74
 blainvillii blainvillii 123
 douglasii hernandesi 177
 m'callii 78, 100
 platyrhinus 78, 100, 123
 solare 177
Phycis chuss 32
 tenuis 32
Phyllodactylus unctus 97
Phyllorhynchus 77
 decurtatus 119, 125
 picta, Chrysemys 37
 Eleotris 68
 picturatus, Hyperolius platyrhinus 179
Pimelometopon pulcher 167
 pinniger, Sebastodes 167
 pipiens, Rana 176
Pituophis catenifer annectens 125
 deserticola 127
 rutilus 177
Planorbis 84
Platax 58
Platynus 102
Platyopocilus 23, 34
 platyrhinus, Phrynosoma 78, 100, 123
 platyrhinus, Hyperolius 179
Plethodon cinereus 85
 glutinosus 85, 102
 wehrlei 95
Pleuronichthys decurrens 67
Pneumatophorus colias 31
 diego 165
 poecilopus, Cottus 17
 poinsettii, Sceloporus torquatus 158
 politus, Seriphus 166
 Spheroides 167
Pollachius virens 32
Polypedates aculeatus 179
 appendiculatus 179
 bimaculatus 179
 eques 179
 goudoti 179
 leucomystax 179
 macrotis 179
 maculatus 179
 mocquardi 179
 ostoni 179
 pardalis 179
 reinwardti 179
 schlegeli 179

- tephraeomystax179
 Pomatomus saltatrix32
 Pomolobus chrysochloris32
 pseudo-harengus32
 Poronotus triacanthus32
 porphyriticus, Gyrinophilus
 8, 82, 83, 85, 101
 Potamophis8
 striatulus7
 princeps, Caulolatilus168
 Prionace glauca31
 Prionotus carolinus32
 proboscideus, Chaenomugil165
 productus, Rhinobatos163
 Profundulus69
 Prosopium62
 coultieri62, 63, 64
 oregonium63
 snyderi62, 63, 64
 williamsoni63
 protenus, Dromicus parvifrons91
 Psammophis72
 Pseudacris triseriata99, 104
 Pseudemys129, 132, 133
 decussata129, 131, 132, 133
 floridana92, 93
 palustris129
 rugosa129, 130, 132, 133
 stejnegeri129, 133
 Pseudobranchius striatus37
 Pseudochromis30
 maccullochi30
 pseudo-harengus, Pomolobus32
 Pseudopleuronectes americanus32
 Pseudotriton duryi81
 montanus81
 ruber ruber101
 pulcher, Pimelometopon167
 pulchra, Anniella4, 5, 6
 punctatus, Chatoessus30
 Diadophis punctatus7, 8
 punctulatus, Hyperolius179
 punctifer, Crotalopsis105
 purpurascens, Nesiotis30
 purpureus, Anoplarchus143
 quadracus, Apeltes33
 radix, Thamnophis113, 114, 116, 117
 Thamnophis radix103
 Raja diaphanes31
 eglanteria31
 erinacea31
 Rana aurora draytonii120
 pipiens176
 sylvatica84
 rasborella, Hemigrammocypris156
 Rasborinus156
 takakii156
 regalis, Cynoscion32
 regilla, Hyla104, 120
 reinwardti, Polypedates179
 erio, Brachydania23
 reticulatus, Acanthophaelus33, 34
 Lebistes33, 147
 Rhinobatos productus163
 Rhinocheilus lecontei126
 rhodoscels, Hyperolius179
 Richmond, Neil101
 riisei, Corynopoma33
 Micurus corallinus7
 roberti, Hyporhamphus164, 180
 robustus, Zoogoneticus69
 roseofusca, Lichanura roseofusca124
 rossii, Hyperolius179
 ruber, Crotalus38
 Pseudotriton ruber101
 rufescens, Chiromantis179
 rufus, Leptopelis179
 rugosa, Emys132
 Pseudemys129, 130, 132, 133
 Runula azalea168
 rutilus, Pituophis catenifer177
 sackii, Cnemidophorus sexlineatus158
 saigonensis, Osphronemus180
 saira, Cololabis66, 67
 salar, Salmo157
 Salmo fario21
 gairdnerii32
 salar157
 saltatrix, Pomatomus32
 Salvadora grahamiae hexalepis124
 santa-anae, Notolepidomyzon136
 Sarda chilensis165
 sarda32
 velox31, 32
 Sardinops caerulea65, 164
 Sargus24
 annulatus24
 Sauromalus obesus121
 Saxidomus giganteus144
 Scaphiopus104
 couchii175
 hammondii36, 104, 119, 175
 holbrookii35
 holbrookii104
 Sceloporus clarkii176
 consobrinus99
 elongatus14
 graciosus graciosus13
 jarrovi177
 magister103, 122
 occidentalis biseriatus122
 torquatus cyanogenys158
 poinsettii158
 undulatus consobrinus158
 schlegeli, Polypedates179
 Schmidt, Karl Patterson6
 Schultz, Leonard P.65, 143
 Scincus73, 76, 77
 officinalis76
 Scleropages30
 formosus30
 Scolecosaurus8
 cuvieri7
 Scomber scombrus31, 180
 Scomberoides156
 Scomberomorus sierra165

- 147
163
126
179
101
33
7
180
69
124
179
38
101
179
132
133
168
177
158
180
67
157
21
32
157
32
124
136
165
32
32
164
24
24
121
144
104
175
175
35
104
176
99
14
13
177
122
122
158
158
158
179
6
143
77
76
30
30
8
7
80
56
65
- scombrus, Scomber31, 180
Scorpius 58
scorpius, Cottus 24
Myoxocephalus 24
scutellatus, Crotalus119, 127
sebae, Ninia atrata 7
Sebastodes atrovirens167
chlorostictus168
constellatus168
elongatus167
goodei167
pinniger167
serranoides167
vexillaris168
secundodorsalis, Thunnus 31
semidoliatus, Geophis7, 8
semifasciata, Triakis163
Seminotus 54
sempervirens, Castanopsis 36
senegalensis, Kassina179
sepoides, Chalcides75, 76, 77, 78
Seriphus politus166
serpentina, Chelydra37
serranoides, Sebastodes167
Serranus21, 25
Sesbania grandiflora 97
setapinnis, Vomer180
Shropshire, Ralph F. 28
sialis, Argentina105
siamensis, Osphronemus180
simonsi, Astroblepus138
sirang, Barbodes 140, 141
Siren lacertina 37
sirtalis, Thamnophis sirtalis...96, 101, 113
Smith, Hobart M.100
Hugh M.156, 180
snyderi, Prosopium 62, 63, 64
solare, Phrynosoma177
Sonora77, 91
occipitalis4, 77, 126
sonorae, Atherinopsis165
soporator, Bathygobius168
Sparus 24
Spheroides annulatus167
maculatus 32
politus167
sphynx, Blennius146
Sphyaena argentea165
borealis 31
Sphyrna zygaena 31
Springer, Stewart105
Squalus acanthias31, 181
Squatina californica 65
stahli, Dromicus 90
stansburiana, Uta121
Uta stansburiana 15
stejnegeri, Pseudemys129, 133
Thamnophis 35
Uta stansburiana122
Stenotomus 24
chrysops22, 23, 24, 25, 32
stephensi, Crotalus confluentus127
Stevardia aliata 33
Stizostedion glaucum182
Stone, Witmer 36
Storer, Tracy I.9, 169
Storeria dekayi 7
victa 6
stouti, Bdellostoma182, 183
striatulus, Potamophis 7
striatus, Centropistes32
Pseudobranchius37
strictogenys, Diadophis punctatus...7, 8
Strongylura exilis164
marina180
subrubrum, Kinosternon subrubrum.. 43
sulcata, Brochymena14
suspectum, Heloderma177
Svihla, Arthur 38
Ruth Dowell 38
Swanson, Gustav 34
sylvatica, Rana 84
Synbranchus 8
marmoratus 7
Synodus sp.164
Syngnathus fuscus180
taeniatum, Arrhyton 90
tagoi, Hynobius158
takakii, Rasborinus156
Tanner, Vasco M.135
Tantilla nigriceps177
Tarpon atlanticus 28
tau, Opsanus 32
Tautoga onitis 32
tenuis, Apostolepis103, 104
Phycis 32
tephraeomystax, Polypedates179
Teratoscincus 73
Terrapene carolina carolina 86
ornata177
tessellatus, Cnemidophorus122
Cnemidophorus tessellatus123
Tetragonopterus viejita137
texana, Holbrookia176
Thamnophis butleri
.....113, 114, 115, 116, 117
megalops 35
ordinoides vagrans99, 103
radix113, 114, 116, 117
butleri 96
radix 103
sauritus proximus103
sirtalis infernalis127
sirtalis96, 101, 113
stejnegeri 35
Thorophos 61
euryops 61
Thunnus secundodorsalis 31
tigrinum, Ambystoma 98
townsendi, Amphisbaena178
Trachidermus ansatus 20
fasciatus17, 20
Trachurops crumenophthalmus165
Tretanorhinus89, 92
triacanthus, Prionotus 32
Triakis semifasciata163
triangulum, Lampropeltis triangulum 98
Trichiurus lepturus180
Trichogaster180

- trichopterus180
 Trimetopon 91
 triseriata, Pseudacris99, 104
 Triturus 1
 viridescens 2
 viridescens 84
 Tropicophis 92
 truei, Ascapus38, 100
 tsuensis, Hynobius158

 Ulca marmorata 67
 ulcerosus, Mantidactylus179
 uluguruensis, Leptophis179
 Ulva180
 Uma73, 75, 76, 77, 78
 notata73, 75, 122
 unctus, Phyllodactylus 97
 undulatus, Menticirrhus167
 Uromacer89, 90, 91, 92
 catesbyi 91
 Uta ornata linearis176
 stansburiana121
 elegans176
 stansburiana 15
 stejnegeri122
 utahensis, Notolepidomyzon135, 136
 utowanae, Anolis 12

 vafer, Myrophis164
 vagrans, Thamnophis ordinoides ..99, 103
 valeriae, Virginia valeriae 7
 valliceps, Bufo158
 Van Hyning, O. C. 37
 variegatus, Coleonyx120
 velox, Sarda31, 32
 ventralis, Callisaurus 72
 Callisaurus ventralis121, 176
 Ophisaurus 7
 vernalis, Liopeltis102
 vexillaris, Sebastodes168
 victa, Storeria 6
 viejita, Tetragonopterus137
 vigilis, Xantusia123
 violaceus, Cebidichthys143
 vipera, Cerastes 73
 virens, Pollachius 32
 virescens, Butoroides 37
 Virginia 8
 valeriae elegans 7
 valeriae 7
 viridescens, Triturus 2
 Triturus viridescens 84
 vittata, Lepidomeda135
 vittatum, Arrhyton 90
 vittatus, Philautus179
 vittiger, Chelonia 14
 Vomer setapinnis180
 vulgaris, Bufo 79
 vulpes, Albula164

 Wales, Joseph H.106, 163
 Walker, Charles F. 81
 wehrlei, Plethodon 95
 Weller, W. Hamilton 81
 whitei, Goodea 68
 williamsoni, Prosopium 63
 willoughbyi, Acrotus 65
 Wilson, Beryl M.169
 wislizenii, Crotaphytus121, 122
 Woodbury, Lowell A. 13

 xanti, Labrisomus168
 Xantusia vigilis123
 xenarcha, Mycteroperca165, 166
 Xenistius californiensis166
 Xenodon 27
 xerampelina, Chiromantis179
 Xiphias gladius 32
 Xystaema cinereum166

 yarrowi, Apocope135
 y-oporus, Leporinus137
 Yucca brevifolia123
 mohavensis123

 Zancus 58
 zonistius, Zoogoneticus 68, 69, 71
 Zoogoneticus68, 69
 cuitzeoensis 69
 diazii 69
 dugesii 69
 maculatus 69
 pachycephalus 69
 robustus 69
 zonistius 68, 69, 71
 zygaena, Sphyrna 31

... 7
... 7
... 2
... 84
... 135
... 90
... 179
... 14
... 180
... 79
... 164

, 163
... 81
... 95
... 81
... 68
... 63
... 65
... 169
... 122
... 13

.168
... 123
... 166
... 166
... 27
... 179
... 32
... 166

.135
... 137
... 123
... 123

. 58
0, 71
3, 69
... 69
... 69
... 69
... 69
... 69
0, 71
... 31